







# DIFFUSE LESIONS OF THE STOMACH

*An Account with Special Reference  
to the Value of Gastric Biopsy*

BY

IAN J WOOD

MD (Melbourne) FRCP FRACP  
Assistant Director of the Walter and Eliza Hall Institute  
of Medical Research Royal Melbourne Hospital and  
Head of the Clinical Research Unit of the Royal  
Melbourne Hospital and the Walter and Eliza Hall Institute

AND

LEON I IATT

MB BS BSc (Melbourne)  
Pathologist to the Clinical Research Unit of the Royal  
Melbourne Hospital and Walter and Eliza Hall Institute  
Melbourne Victoria Australia



LONDON

EDWARD ARNOLD (PUBLISHERS) LTD

© I J Wood and L I Taft 1958

*First published 1958*

Made and printed in Great Britain by  
William Clowes and Sons Limited London and Beccles

TO  
*F M BURNET, F R S*

WHO OFTEN TOLD US THAT  
NEW KNOWLEDGE FOLLOWED CLOSELY  
IN THE WAKE OF A  
NEW TECHNIQUE



# CONTENTS

<i>Chapter</i>		<i>Page</i>
1	<b>METHODS OF INVESTIGATION</b>	1
	Problems of the early investigators	1
	Symptomatology and clinical investigation	2
	Test meal examination	3
	Radiology	4
	Gastroscopy	5
	Gastric biopsy	6
	Technique of gastric biopsy with the flexible gastric biopsy tube	7
	Preparation and staining of biopsy specimens for microscopic examination	11
	The appearance of the normal gastric biopsy	13
	The indications for gastric biopsy using the flexible tube	15
	Complications of gastric biopsy with the flexible tube	16
	Gastric cytology	17
2	<b>ACUTE GASTRITIS</b>	18
	Structural changes	18
	Symptomatology	20
	Aetiology	22
	Treatment	23
3	<b>CHRONIC GASTRITIS</b>	24
	Structural changes	24
	Symptomatology of chronic atrophic gastritis	33
	Aetiology of chronic atrophic gastritis	37
	Relationship of chronic gastritis to peptic ulcer	45
	Relationship of chronic gastritis to carcinoma	46
	Relationship of chronic gastritis to anaemia	48
	Treatment	51



<i>Chapter</i>		<i>Page</i>
4	DIFFUSE GIANT HYPERTROPHIC GASTRITIS	53
5	GASTRIC ATROPHY PERNICIOUS ANAEMIA AND SUBACUTE COMBINED DEGENERATION OF THE CORD	57
	Structural changes of gastric atrophy and per- nicious anaemia	57
	Symptomatology of gastric atrophy and pernicious anaemia	61
	Aetiology of gastric atrophy and pernicious anaemia	64
	The relationship of the gastric changes to pernicious anaemia subacute combined degeneration of the cord and other megaloblastic anaemias	66
	Can pernicious anaemia be diagnosed in the pre- sence of normal acid secretion and normal appearance of the gastric biopsy ?	68
	Definition of pernicious anaemia	69
	Treatment of gastric atrophy pernicious anaemia and subacute combined degeneration of the cord	69
	EPILOGUE	72
	REFERENCES	73
	INDEX	82

## PREFACE

In 1946 our Clinical Research Unit was formed by Sir Macfarlane Burnet F R S Director of the Walter and Eliza Hall Institute to work in the Institute and in the wards of the Royal Melbourne Hospital. Diseases of the gastro intestinal tract were chosen for study and we soon encountered middle aged or elderly patients who had suffered from flatulent dyspepsia for many years. They exhibited diffuse epigastric tenderness and an absence of the usual causes of dyspepsia such as ulcer cancer gallstones and pancreatitis. We first studied the gastric secretions and found them to be depleted and in 1948 we developed the technique of gastric biopsy using a flexible gastric biopsy tube designed in the unit (Wood Doig Motteram and Hughes 1949).

Now some 10 years after our study began we are privileged to present our collected observations in this book. Much of the data has been selected from our published papers which have appeared in Australia England and the United States of America. We have added a limited number of observations made by other workers in this field many of whom have generously helped us over the years. Our special thanks are due to Dr Avery Jones and Professor Magnus in England Dr Walter Palmer and Lieutenant Colonel Eddy Palmer in the United States of America and in Australia to members of the Medical Staff of the Royal Melbourne Hospital particularly Dr W E King Mr J O Smith Mr Grayton Brown Dr E Graeme Robertson Dr D C Cowling and finally and most gratefully to our many colleagues who have worked in our Clinical Research Unit over the years particularly Dr R Doig Dr S Weiden Dr R Motteram Dr J Funder Dr E Finckh Dr R Joske and Dr I Mackay together with their technical assistants Miss E Davis and Miss E Earle. Mr E Matthaer of the Melbourne University and Mr R Ingles of the Royal Melbourne Hospital kindly helped with the photography.

The title *Diffuse Lesions of the Stomach* has been used to include diffuse gastritis in its acute and chronic forms gastric atrophy and that rare disease diffuse giant hypertrophic gastritis. It naturally includes pernicious anaemia and excludes the localised lesions of

peptic ulcer and cancer although some reference is made to their relationship to gastritis

All the studies have been undertaken in adults as no children are treated in the Hospital It may well be that some of the lesions seen in adult life have begun during childhood or even in the prenatal period

## CHAPTER 1

# METHODS OF INVESTIGATION

### Problems of the early investigators

For many centuries physicians considered dyspepsia was frequently caused by diffuse inflammation of the stomach. The celebrated Broussais (1831) initiated widespread belief in the occurrence of chronic phlegmasiae or inflammations of the stomach by reporting observations which he made while military surgeon to the army of Napoleon.

However many factors confused early investigators in this field especially their failure to appreciate the gross changes which took place in the gastric mucosa soon after death. In later years recourse was made to the study of specimens obtained at surgical operation but even here considerable changes were found to be brought about by the manipulations by the surgeon. An appreciation of these problems created a wave of disbelief in the prevalence of gastritis especially in its chronic stage. Moreover even when physicians did accept its presence they were loath to accept it as a cause of chronic dyspepsia.

A challenge to this disbelief came at the turn of the present century led by the splendid studies of Faber (1935) who prevented post mortem digestion of the gastric mucosa by introducing formalin immediately after death. But this post mortem material could still be criticised on the grounds that structural abnormalities might first appear during the terminal illness (see Chapter 3).

There was an urgent need for fresh undamaged specimens of gastric mucosa which could be placed immediately in a fixative solution and prepared for histological examination. Moreover it would be of considerable value if these specimens were obtained with minimal discomfort and risk to the patient for this would enable a comprehensive study to be made not only in health and disease but also over a number of years in the same individual.

*It is our sincere belief that the technique of gastric biopsy first*

described in our Unit in 1949 has made a liberal contribution to this urgent need

In this chapter the methods used in our studies are discussed

## Symptomatology and clinical investigation

In addition to the history and physical examination it may be necessary to employ a fractional test meal X ray by barium meal and even gastroscopy and gastric biopsy to establish the diagnosis of gastritis and exclude other diseases with similar clinical features

*Clinical history* The taking of a comprehensive history is of considerable importance It may call for much time and patience from the physician but the information is rewarding The family history social history and the nature of past illnesses may be of value

In taking the history and making the physical examination it should be constantly recalled that diffuse lesions of the stomach have to be differentiated from gastric and duodenal ulcer cancer of the stomach chronic pancreatitis and other diseases involving the upper portion of the abdomen Pure functional dyspepsia will always present a problem in diagnosis often true gastritis is present with considerable functional overlay

*Pain and discomfort* The nature site and duration of the discomfort should be ascertained together with its provocation by food physical exertion or mental stress The type of food which causes pain is of interest—pickles pastry and sherry are often incriminated The relief of pain by alkalis dilute hydrochloric acid or food should be ascertained

*Diet* The appetite and the quality and quantity of the food consumed should be recorded together with the general eating habits In Australia as indeed in many other parts of the world the consumption of alcohol is often excessive and this may cause gastric disturbances due either to its direct effect on the gastric mucosa or to its influence on the diet

*Physical examination* In carrying out the general physical examination the local signs of gastric disease should be noted and also a special search should be made for lesions elsewhere in the body which may be associated with gastritis as they may share in a common aetiology Thus hepatomegaly and peripheral neuritis may suggest alcoholism and malnutrition

There may be premature greying of the hair atrophy of the tongue recurrent stomatitis haemorrhagic gums carious teeth or complete dental extraction anaemia purpura pigmentation or vascular spiders in the skin cardiac enlargement and tachycardia from vitamin deficiency and finally peripheral neuritis or early signs of subacute combined degeneration of the cord

Examination of the upper abdomen may reveal the diffuse epigastric tenderness of gastritis in contrast to the more local tenderness of peptic ulcer gall bladder disease or perhaps cancer. A mass in the epigastrium may be hepatic enlargement hydatid disease or cancer of the stomach. Rarely it may be a tumour of the body of the pancreas or enlargement of the retroperitoneal glands. A more laterally placed mass may be caused by a palpable gall bladder splenic enlargement or a renal tumour

### Test meal examination

The histamine test meal is a most satisfactory method of determining gastric secretion and thereby obtaining some evidence as to the state of the gastric mucosa. There is good correlation between the test meal and the findings in gastric biopsy as is clearly shown in fig 1 (Wood *et al* 1949 Joske *et al* 1955) (see Chapter 5)




Fig 1 The relationship between the findings on gastric biopsy and the histamine test meal. With the increase in the inflammatory reaction and atrophy of the gastric mucosa there is a corresponding fall in the secretion of acid. There were 275 biopsies within normal limits 296 showing superficial gastritis 172 atrophic gastritis and 41 gastric atrophy. (Data from Joske Finckh and Wood 1955)

In most of our series a standard subcutaneous dose of 0.9 mg of histamine acid phosphate was used. This stimulus produced a satisfactory rise in acid in normal controls and high levels in cases

of duodenal ulcer, but without producing unpleasant histamine effects in the patient. Before giving histamine the stomach was emptied by passing a fine stomach tube such as the Rehfuß model, and then after giving histamine it was emptied at half hourly intervals for one and a half hours. In some cases of suspected pernicious anaemia the maximum secretory response was evoked by Hay's method (1953) when an antihistamine drug was given to counter the general systemic effects of a liberal dose of histamine (0.4 mg histamine acid phosphate per 10 kg body weight) as the antihistamine drugs do not inhibit the histamine stimulus to gastric secretion.

Hypoglycaemia caused by intravenous insulin stimulates gastric secretion by way of the vagus nerve and an insulin test meal (insulin 15 units intravenously) is of value after vagotomy to assess the degree of completeness of the nerve section.

An overnight test meal reveals the resting secretion over a long period continuous gentle suction or 2 hourly aspiration being used.

The tubeless test meal in which the excretion of uropepsin is estimated after histamine stimulation has not been used in this series as it is less accurate than the direct measurement of gastric secretion (Levy and Levine 1956) however it is of value as a screening test for lowered gastric secretion. The plasma pepsinogen level may be a guide to gastric secretion. Hoar and Browning (1956) claim that it is superior to urinary pepsinogen determination.

A tubeless test meal using cation exchange resins may also be used as a screening test for anacidity (Rechtschaffen and Weingarten 1957).

## Radiology

The radiological examination is of considerable value in the diagnosis of localised lesions of the stomach and duodenum notably peptic ulcer, cancer, diverticulum and prolapse of gastric mucosa through the pylorus. However the value of radiology in detecting diffuse lesions of the stomach is mostly confined to the rare disease diffuse giant hypertrophic gastritis (see Chapter 4). It is of little value as a method of detecting chronic atrophic gastritis or gastric atrophy (Joske *et al* 1955). Furthermore in our experience the diagnosis of hypertrophic gastritis based on the findings of pronounced gastric rugae has been proved by

biopsy to be incorrect except in diffuse giant hypertrophic gastritis. After considerable experience Eddy Palmer (1954) has reached a somewhat similar opinion to that held in our unit.

### Gastroscopy

Most patients in our series were examined by gastroscopy. The technique consists of spraying the throat with local anaesthetic (cocaine 1.5 ml of 3 per cent with adrenalin 1:10 000) preceded by basal sedation with chlorpromazine 50 mg, omnopon 20 mg and hyoscine 0.4 mg. The stomach is emptied by gravity using a large bore stomach tube and elevating the foot of the examination couch.

The Hermon Taylor gastroscope was used in 95 per cent of our last 200 gastroscopies as it has a controllable lower segment which affords a good view of most localised lesions. However in elderly patients especially in females over the age of 60 the less bulky Cameron model (Schindler design) was used to reduce the hazard of pharyngeal or oesophageal perforation. We encountered only one such perforation in 661 gastroscopies. The patient recovered after drainage of a paraoesophageal abscess which developed in spite of vigorous treatment with antibiotics. Katz *et al* (1956) described a massive pneumoperitoneum after gastroscopy. This subsided with conservative treatment.

Gastroscopy may show congestion, mucopurulent exudate and some petechial haemorrhages and even erosions in acute gastritis and during the acute relapses in chronic gastritis. In pernicious anaemia with gastric atrophy the atrophy may be observed in the body of the stomach especially in the upper segments where the underlying veins are revealed.

However gastric biopsy has provided evidence that gastroscopy often fails to determine whether chronic gastritis is present. Thus Joske *et al* (1955) reviewed 258 instances where gastric biopsy had been performed within 2 weeks of the gastroscopy, often at the same time. In 164 instances where gastroscopy reported the gastric mucosa to be normal, gastric biopsy revealed normal gastric mucosa in only 87 instances, whereas superficial gastritis was present in 56, atrophic gastritis in 16 and gastric atrophy in 5. Although these findings could be partly due to a sampling error in the gastric biopsies, the study does emphasise the limitations of gastroscopy as a method of diagnosing chronic gastritis.



## Gastric biopsy

In 1948 one of us (I J W) observed that fragments of gastric mucosa of sufficient size for microscopic examination might be obtained during a histamine test meal when vigorous suction was exerted through the stomach tube during the collection of the gastric juice. Moreover it was found that better fragments were obtained when negative pressure was exerted and then the tube was suddenly jerked out so as to dislodge a fragment. These fragments could be fixed immediately in formalin embedded in paraffin cut and stained. However they contained mostly superficial epithelium.

Our initial observation was subsequently found to have been recorded by Einhorn in 1894 and later by Hawksley in 1939 who subsequently heard of Einhorn's classical observation (Hawksley and Cooray 1948).

*The development of the flexible gastric biopsy tube.* Our findings of such fragments stimulated the development in our unit of the flexible gastric biopsy tube which now has been used by the unit in carrying out 1736 successful biopsies. It has also been used in its present or a modified form by other workers in many parts of the world notably by E. D. Palmer (1950) in the United States of America who has made many excellent contributions to the knowledge of gastric histopathology.

Others who have made valuable studies with this biopsy technique either in the form described or with a modified model include Badenoch and Richards (1953) Rubins *et al* (1953) Coghill and Williams (1955) Edwards and Edwards (1956) Markson and Davidson (1956) and Shiner and Doniach (1957).

*Chevalier Jackson and gastric biopsy.* The credit must be given to Chevalier Jackson the doyen of endoscopists for first using gastric biopsy under direct vision and establishing it as a valuable and safe method of investigating diseases of the stomach. In 1906 he first performed gastric biopsy using forceps through his rigid oesophagoscope and later stated "In malignant diseases a specimen may be taken with little risk in fungating conditions but in flat ulcerations suspected of malignancy the biting out of the edge of the ulcer though very easy of accomplishment is unjustifiable" (Jackson 1907). His work proved of considerable value and in 1935 he and his son reported "The taking of tissue through the open tube gastroscope is easily done and hundreds of cases without a single complication have demonstrated its freedom from

danger when done with proper precautions (Jackson and Jackson 1935)

In 1937 Schindler described the flexible gastroscope a momentous discovery in the field of gastroenterology. Later Hermon Taylor designed a gastroscope with a controllable lower segment which afforded the operator a more extensive view of the stomach. Schindler's discovery led to the development of the operating gastroscope containing a channel through which forceps can be manipulated under direct vision. Now many but by no means all of the technical difficulties have been overcome following the researches in the United States of America by Kenamore (1940), Kenamore, Scheff and Womack (1946), Benedict (1948) and Tomenius (1952).

Most of our studies described in this book were made with our flexible gastric biopsy tube, the Hermon Taylor gastroscope or the Benedict operating gastroscope.

Each instrument has its own particular virtues and limitations. For diffuse lesions of the stomach the flexible gastric biopsy tube obtains one or two satisfactory fragments of gastric mucosa with minimal inconvenience to the patient. In contrast to this the Benedict operating gastroscope is more bulky and may cause considerable discomfort during and after the examination. However it enables fragments of mucosa to be obtained under direct vision from most parts of the stomach although fragments from antral lesions are difficult to obtain. This type of instrument is thus essential for obtaining tissue from local lesions such as ulcer or carcinoma. We have used the Benedict operating gastroscope on 53 occasions after preliminary inspection with the Hermon Taylor instrument. Successful biopsies were obtained on 40 occasions and there were no complications.

### Technique of gastric biopsy with the flexible gastric biopsy tube

The technique of gastric biopsy using the flexible gastric biopsy tube designed in our unit is described in detail as this method is most suitable for studying diffuse lesions of the stomach. This procedure has been the *piece de resistance* of our work.

The examination can be made on out patients so long as they are not unduly debilitated or anaemic.




Fig. 2. Flexible gastric biopsy tube.

*Top illustration* — The complete biopsy tube with operating handle and wire (top) length of tube with white marks at levels I, II and III and the operating head (bottom) (Bv. 1)

*Center illustration* — Operating head with terminal screw (left) and cylindrical blade to be thrust down and unscrewed from the operating wire. This enables the biopsy fragment to be obtained (x 14)

*Lower illustration* — Operating head assembled. The cutting edge of the cylindrical knife blade is visible through the lateral hole (x 14)

biopsy tube consists of a length of coiled Bowden wire covered by plastic tubing to render it airtight (fig 2) (Wood *et al* 1949<sup>1</sup> and II Doig and Wood 1952)\* At the lower end a steel cylinder is attached which houses a cylindrical knife blade with a proximal cutting edge The end of the cylinder is closed with a terminal screw and its side is pierced by a countersunk circular lateral hole  $\frac{3}{16}$  inch in diameter through which a fragment of gastric mucosa may be sucked The fragment is then cut off by pulling up the cylindrical knife past the lateral hole by means of a length of wire which traverses the tube to emerge through a vacuum tight gland at the proximal end Suction is applied with a pump through a lateral tube which emerges close to the proximal end The overall length of the biopsy tube is 100 centimetres and levels I II and III are marked 45 55 and 65 cm respectively from the lateral hole in the operating head

*Technique* The patient is examined fasting and no sedative is given The pharynx is anaesthetised by a gargle with 15 ml of 3 per cent cocaine with adrenalin 1:10 000 A fine stomach tube is passed into the stomach with the patient sitting The stomach is then completely emptied by the patient lying prone on a couch which is inclined to an angle of 30° with the head downmost and by applying suction to the tube with a 20 c.c. syringe while it is slowly withdrawn Should air or fluid remain in the stomach the chance of obtaining a fragment of mucosa is greatly diminished

The flexible gastric biopsy tube is then passed with the patient lying on his left side the passage through the pharynx being aided by vigorous swallowing Rarely the tip is arrested by spasm at the cardia and this can usually be overcome by encouraging deep breathing and swallowing If still arrested it may pass if the patient sits erect or it may be necessary to administer an inhalation of amyl nitrite

The tube is introduced past mark I till mark II approaches the lips The fragment is then cut off by (a) forcing down the blade by means of the control wire to open the lateral hole (b) immediately exerting negative pressure by means of the exhaust pump to suck in the fragment (c) firmly pulling up the blade

The tube is then withdrawn 2-4 cm and rotated 180° in order to engage a new area of gastric mucosa and the cutting is repeated to obtain a second fragment

\* This flexible gastric biopsy tube is manufactured by Drug Houses of Australia Ltd Alfred Place Melbourne Victoria Australia

The tube is removed for inspection and the patient instructed to lie still. The fragments of gastric mucosa are located by removing the terminal screw forcing out the cylindrical knife blade and unscrewing the blade. The fragments will be found either inside the cylindrical knife blade or more frequently within the steel cylinder which houses the blade. They are dislodged from the cylinder by directing a current of air down the tube.

The fragments are immediately fixed in 10 per cent formol saline thus providing excellent material for histological examination. The instrument is immediately cleaned by flushing it through with a liberal quantity of water and then drying with a stream of air. The tube can then be sterilised with Zephiran 1:100 and the knife blade with alcohol. The instrument is then placed in a sterile cloth bag. It is essential that the knife blade be sharpened after every five operations.

After the biopsy has been performed the patient has complete rest lying down for a period of 4 hours. Sips of milk are given after 1 hour and a light meal at the end of 4 hours. Out patients are then sent home to rest arrangements being made for them to be driven home. They return to full work on the following day but within 48 hours they visit the Hospital for a medical inspection to exclude haemorrhage.

In our cases gastric biopsy was usually a painless procedure—on rare occasions a minor dull burning sensation was experienced but this was usually caused by vigorously thrusting the tube too far into the stomach thus exerting pressure on the greater curvature for it was immediately relieved by withdrawing the instrument 2–3 cm. After the operation was completed a few patients had minor epigastric discomfort for 12–24 hours but the significance of this was difficult to assess as most of them were being investigated for a similar complaint.

In general it may be said that gastric biopsy performed with the flexible gastric biopsy tube is not a major undertaking. Most patients are quite prepared to have it repeated at intervals of 1 or 2 years.

*The efficiency of gastric biopsy with the flexible tube.* The efficiency of this method is shown by the combined experience of a number of operators in our unit. Between the years 1946 and 1957 there were 1949 attempts at gastric biopsy with the flexible tube and 89 per cent were successful. The unsuccessful attempts were due to failure to obtain a suitable fragment in 6 per cent and the

obtaining of oesophageal mucosa in 5 per cent. The 1736 successful biopsies were carried out on 1046 patients many patients having more than one biopsy. Thus in one patient 15 successful biopsies were carried out over a period of 8 years.

**Sampling error** As the area of mucosa examined in a single biopsy is so small it is inevitable that a sampling error occurs. To minimise and measure this error usually 2 fragments of mucosa are obtained from different portions of the body of the stomach with each passage of the biopsy tube. Joske *et al* (1955) found that of 726 cases in 73.8 per cent the appearances in the 2 fragments were similar while in 26.2 per cent there were sufficient differences between the fragments to suggest that the changes were patchy. Similarly of 123 patients in whom serial gastric biopsies were carried out 76.4 per cent showed substantial agreement in histology while 23.6 per cent did not. Some of the latter were due to known local lesions (ulcer carcinoma) and in others they were related to the changing course of the illness.

Studies of operation and autopsy material reveal that gastritis usually involves the body of the stomach fairly diffusely (Hebbel 1949 Williams 1950). Furthermore it has been found that the results of the histamine test meal and the biopsy appearance are related in that the lower the gland content of the section the lower the acidity and output of the gastric juice (Funder and Weiden 1952).

### **Preparation and staining of biopsy specimens for microscopic examination**

The fragment of gastric mucosa obtained by the flexible biopsy tube consists of a disc of tissue approximately 3 mm in diameter and 0.5 mm in depth (fig. 3). Immediate fixation is carried out in 10 per cent formal saline for 6 to 12 hours. It is then prepared for paraffin embedding in the routine manner.

It is essential to obtain vertical sections of mucosa. Should difficulty be experienced in achieving this the fixed specimen may be cut perpendicularly to the surface with a sharp scalpel and appropriately placed in the molten paraffin during embedding. Sections 5  $\mu$  in thickness are cut, mounted and stained with eosin and haematoxylin, haematoxylin and Mayer's mucicarmine for detecting alterations in the mucus secreting cells and the trichrome stain of Mottram (1951) which is a modification of that

used by Bowie and Vineberg (1935) The trichrome stain is most informative as it clearly identifies the chief and parietal cells It is used as follows



Fig 3 A small gastric biopsy fragment obtained with the flexible gastric biopsy tube to show the specimen of mucosa obtained (eosin and hematoxylin  $\times 45$ )

#### *Reagents for trichrome stain*

##### 1 Pepsinogen stain Stock solution

Methyl violet 6B (1 per cent in aqueous solution)—100 ml

Orange G (0.5 per cent in aqueous solution)—100 ml

Mix allow to stand overnight centrifuge and wash the precipitate 6 times Dissolve the precipitate in 100 ml absolute ethyl alcohol

For use take 2 parts of the stock solution and add one part of distilled water The stock solution is quite stable in a well stoppered bottle

##### 2 Ponceau—fuchsin—phloxine

Ponceau de xyline (1 per cent in aqueous solution)—6 ml

Acid fuchsin (1 per cent in aqueous solution)—2 ml

Azophloxine (0.5 per cent in aqueous solution)—2 ml  
 Acetic acid (1 ml glacial acetic acid in 1500 ml water)—88 ml

### 3 Methyl green (1 per cent in aqueous solution)

#### *Staining method for trichrome stain*

- 1 Pepsinogen stain—10 minutes in a Coplin jar
- 2 Decolourise with 70 per cent ethyl alcohol until the violet colour remains in the chief cells only
- 3 Wash with water
- 4 Ponceau—fuchsin—phloxine—10 seconds
- 5 Wash with water
- 6 Methyl green—60 seconds
- 7 Differentiate rapidly with 95 per cent ethyl alcohol
- 8 Dehydrate with absolute alcohol
- 9 Xylol
- 10 Mount in Canada balsam

*Results of trichrome stain* Pepsinogen granules purple  
 parietal cell cytoplasm red Paneth cell granules dark red  
 nuclei bluish green goblet cell mucus green

### **The appearance of the normal gastric biopsy**

Detailed accounts of gastric cytology have been given by Bensley (1928) Plenck (1932) and Maximow and Bloom (1952). Microscopic examination of biopsy fragments from the body of the normal stomach shows the mucosa to consist of a dense aggregation of regular branched tubular glands opening through a neck into the *foveolae gastricae* or gastric pits (fig. 4). The epithelium of the pits and glands is supported by a basement membrane formed by a condensation of the *lamina propria*; the glands extending down to the *muscularis mucosae*. The average depth of the processed section from the surface to the inner aspect of the *muscularis mucosae* is 0.54 mm (Joske *et al.* 1955).

The pits extend down into the mucosa for a distance of one third to half of its depth and their lumen (10–40  $\mu$ ) may be wider than that of the glands. The superficial epithelium lining the surface of the mucosa and extending into the pits consists of a single layer of regular tall columnar epithelium with small round or oval basal nuclei and a pale eosinophilic cytoplasm containing a theca of mucigenic granules which stain with mucicarmine. Goblet cells are not present in the normal gastric mucosa. Smaller mucigenic cells are also present in the necks of the glands. Mucus



may be visible in the lumen of the pits forming an adherent layer on the superficial epithelium

Bensley (1928) found mitoses in the cells lining the bases of the pits so that replacement of the rapidly shedding gastric epithelium probably occurs by multiplication of the cells in these areas. This is borne out by the progressive differentiation in the cells from the depths of the pits to the surface, sometimes becoming more pronounced during recovery from injury (Motteram 1951). We have not encountered obvious mitoses in the biopsy specimens prepared by our technique but E. Palmer (1954) has reported an excellent example in a case of acute alcoholic gastritis.

The parietal or acid secreting cells are more numerous in the superficial portion of the glands polyhedral in shape with the deeper cells tending to be displaced laterally from the lumen of the gland. Their cytoplasm is eosinophilic. The pepsinogen secreting or chief cells are more numerous in the deeper portion of the glands cuboidal or low columnar in shape with basal nuclei and granular cytoplasm. Argentaffine cells are not seen in sections stained in the manner described above.

In the antrum and pyloric areas the pits are deeper and the glands here consist of coiled tubules lined by low columnar or cuboidal cells with basal nuclei and a pale granular cytoplasm which stains with mucicarmine. They resemble the Brunner's glands of the duodenum. Interstitial cellularity is more pronounced. The glands in the region of the cardia are similar in appearance to the glands of the antrum.

The *lamina propria* is a delicate reticulum containing a few fibroblasts, plasma cells and lymphocytes. Capillaries and lymphatics in this area are occasionally apparent in the sections. Occasionally a small lymph follicle is present in the deeper portion of the mucosa. In the antrum interstitial cellular infiltration is more pronounced. The *muscularis mucosae* forms a lamina at the base of the glands sending an occasional fibre up into the *lamina propria* between the glands. In the biopsy the circular and longitudinal fibres are not seen as such because of the random orientation of the fragment. In some biopsies the connective tissue and small vessels of the submucosa can be seen.

Artefacts due to the suction of the biopsy cylinder include small extravasations of red cells under the superficial epithelium and the aspiration of epithelial or gland cells into the lumen of the glands or pits. In abnormal specimens both these phenomena are more



Fig 4 Normal gastric mucosa in a gastric biopsy. With the trichrome stain the parietal (red) and chief cells (purple) are easily distinguished in the glands (Trichrome stain  $\times 80$ )



Fig 5 Chronic atrophic gastritis with moderate atrophy. Patchy intestinal metaplasia with goblet cells in the superficial epithelium. The gastric pits are distorted, the glands are reduced in number and contain non-specific cells. The lamina propria is infiltrated with inflammatory cells (Trichrome stain  $\times 80$ )




Fig 6 Gastric atrophy. The biopsy shows diffuse intestinal metaplasia and complete atrophy of the glands (Trichrome stain  $\times 110$ )

may be visible in the lumen of the pits forming an adherent layer on the superficial epithelium

Bensley (1928) found mitoses in the cells lining the bases of the pits so that replacement of the rapidly shedding gastric epithelium probably occurs by multiplication of the cells in these areas. This is borne out by the progressive differentiation in the cells from the depths of the pits to the surface, sometimes becoming more pronounced during recovery from injury (Motteram 1951). We have not encountered obvious mitoses in the biopsy specimens prepared by our technique but E. Palmer (1954) has reported an excellent example in a case of acute alcoholic gastritis.

The parietal or acid secreting cells are more numerous in the superficial portion of the glands polyhedral in shape with the deeper cells tending to be displaced laterally from the lumen of the gland. Their cytoplasm is eosinophilic. The pepsinogen secreting or chief cells are more numerous in the deeper portion of the glands cuboidal or low columnar in shape with basal nuclei and granular cytoplasm. Argentaffine cells are not seen in sections stained in the manner described above.

In the antrum and pyloric areas the pits are deeper and the glands here consist of coiled tubules lined by low columnar or cuboidal cells with basal nuclei and a pale granular cytoplasm which stains with mucicarmine. They resemble the Brunner's glands of the duodenum. Interstitial cellularity is more pronounced. The glands in the region of the cardia are similar in appearance to the glands of the antrum.

The *lamina propria* is a delicate reticulum containing a few fibroblasts plasma cells and lymphocytes. Capillaries and lymphatics in this area are occasionally apparent in the sections. Occasionally a small lymph follicle is present in the deeper portion of the mucosa. In the antrum interstitial cellular infiltration is more pronounced. The *muscularis mucosae* forms a lamina at the base of the glands sending an occasional fibre up into the *lamina propria* between the glands. In the biopsy the circular and longitudinal fibres are not seen as such because of the random orientation of the fragment. In some biopsies the connective tissue and small vessels of the submucosa can be seen.


Artefacts due to the suction of the biopsy cylinder include small extravasations of red cells under the superficial epithelium and the aspiration of epithelial or gland cells into the lumen of the glands or pits. In abnormal specimens both these phenomena are more



Fig 4 Normal gastric mucosa in a gastric biopsy. With the trichrome stain the parietal (red) and chief cells (purple) are easily distinguished in the glands (Trichrome stain  $\times 80$ )



Fig 5 Chronic atrophic gastritis with moderate atrophy. Patchy intestinal metaplasia with goblet cells in the superficial epithelium. The gastric pits are distorted, the glands are reduced in number and contain non-specific cells. The lamina propria is infiltrated with inflammatory cells (Trichrome stain  $\times 80$ )



Gastric atrophy. The biopsy shows diffuse intestinal metaplasia and complete atrophy of the glands (Trichrome stain  $\times 110$ )



pronounced and may reflect actual changes of this nature or the more ready production of such changes in the abnormal mucosa

### The indications for gastric biopsy using the flexible tube

Gastric biopsy is of considerable value in determining the cause of chronic dyspepsia where neither X ray nor gastroscopy reveals an ulcer or cancer. Here a biopsy may aid in the diagnosis by showing active chronic gastritis with considerable atrophy of the parietal and chief cells. Active chronic gastritis may be the sole cause of the symptoms and signs. It must be appreciated however that the biopsy *per se* does not exclude cancer.

Biopsy may also indicate gastritis as being the cause of chronic iron deficiency anaemia from continuing blood loss. This blood loss is usually in minimal quantity but from time to time there may be a frank melæna and even a hæmatemesis.

Gastric biopsy may also aid in the diagnosis of subacute combined degeneration of the cord where examination of the peripheral blood and bone marrow reveals no clear evidence of pernicious anaemia and where the estimation of serum vitamin B<sub>12</sub> is not available or is rendered valueless by previous injections of vitamin B<sub>12</sub>. The more elaborate absorption tests following the oral administration of <sup>57</sup>cobalt labelled vitamin B<sub>12</sub> is also informative in these circumstances (Booth and Mollin 1956; Witts 1956).

Gastric biopsy is contra indicated when there is clinical evidence of a bleeding tendency supported by laboratory tests. Also neither gravely ill patients nor patients with severe anaemia should be examined in this way—the latter should first be treated by transfusion or other suitable means to relieve their anaemia. This applies to cases of megaloblastic anaemia where the cause is in doubt. Should it be true pernicious anaemia the gastric atrophy will remain unchanged in spite of full treatment with vitamin B<sub>12</sub> (Finckh and Wood 1953).

Gastric biopsy will also aid in determining the cause of achlorhydria for there is a close correlation between the gastric biopsy findings and the histamine test meal (fig. 1). This was first shown by Wood, Doug, Weiden and Moore (1949) and more recently by Shiner and Doniach (1957) who found good correlation between gastric biopsy and the secretion of acid and pepsin.

### Complications of gastric biopsy with the flexible tube

Haemorrhage into the lumen of the stomach was the only complication encountered in our series. There was no evidence of gastric perforation—the instrument was designed to cut down to the *muscularis mucosae* and no deeper and this was confirmed by extensive preliminary trials on the bench using fresh specimens of stomach removed at operation. Gastroscopy performed immediately after biopsy revealed a minute lesion usually covered by blood clot, at the most there was minimal bleeding. There was no evidence of persistent ulceration as judged by the absence of pain. X ray examination gastroscopy or examination of the mucosa when subtotal gastrectomy for duodenal ulcer was performed one week or more after the biopsy was taken. However a small superficial healing lesion could usually be observed in a stomach resected 1 or 2 days after the biopsy.

Using the same model of the flexible gastric biopsy tube and working with cats Gunter (1950) made detailed observations on the rate of healing of the gastric mucosa after biopsy and found that a single layer of regenerating epithelium just covered the lesion in 3 days and experience with humans indicated a similar rate of healing.

Haemorrhage sufficient to cause symptoms of anaemia haematemesis or melaena or a pronounced fall in haemoglobin occurred in only 10 of 1947 attempted biopsies. 87 per cent of the latter providing suitable fragments of gastric mucosa.

Most haemorrhages occurred within 12 hours of the biopsy and caused weakness sweating and pallor followed by the passage of a melaena stool. Rarely there was a small haematemesis. Limited blood transfusion was required in only 2 of the 10 patients. 1 of these was severely and chronically ill with subacute combined degeneration of the cord and bladder sepsis. There were no fatalities.

In our series haemorrhage has been much less frequent since patients have been kept at strict rest in bed for 4 hours after making the biopsy and since biopsy has not been undertaken in patients showing the bleeding diathesis.

Markson and Davidson (1956) encountered 2 cases of haematemesis in 111 gastric biopsies performed with the flexible tube one an elderly male with hypertension and renal insufficiency the other a middle aged female with untreated pernicious anaemia. Both responded to blood transfusion. Using a modification of the

flexible tube Rubin *et al* (1953) reported haemorrhage in 2 of over 200 biopsies and Coghill and Williams (1955) recorded haematemesis in 9 (2.1 per cent) of 1189 attempted biopsies on 437 patients and also one case of perforation of the oesophagus. E. Palmer and Smith (1953) reported 2 patients who developed a submucosal haematoma but recovered with conservative treatment.

Shiner and Doniach (1957) performed 127 successful biopsies without any complications and there were few failures to obtain a suitable fragment.

The best results are obtained by gentle handling of the instrument till it is in position by keeping the head of the instrument stationary while the negative pressure is applied and the knife blade firmly pulled up ensuring that the knife is sharpened frequently and arranging that after the biopsy the patient has absolute rest for 4 hours. And finally, gravely ill patients and patients with a bleeding tendency should not be subjected to biopsy.

Such measures should reduce the complications to less than 1 per cent. Severe complications should be rarely encountered.

### Gastric cytology

Because of our interest in gastric biopsy we have only had limited experience with the study of gastric washings. Undoubtedly this method is of value in the detection of gastric cancer and even pernicious anaemia when it is performed by pathologists skilled in this procedure (Massey and Rubin 1954; Crozier *et al* 1956).



## CHAPTER 2

# ACUTE GASTRITIS

### Structural changes

Surely the most dramatic accounts of acute inflammation in the stomach have come from Beaumont (1833) who described the hyperaemia mucus patches erosions and diminution in secretion in the gastric fistula of Alexis St Martin following strong alcoholic beverages and other dietary trauma and during acute febrile episodes. These studies have been paralleled in the modern era by Wolf and Wolff (1943) who demonstrated that psychic stimuli in their patient Tom produced acute inflammatory changes in the mucosa of his gastric fistula. Gastrosopic appearances in acute gastritis have been recorded by Schindler (1922) and E Palmer (1951) while haematogenous gastritis in fatal infections has been found by Vimtrup (1929) and Nyfeldt and Vimtrup (193-) using the technique of Faber (1935) of early post mortem fixation of the stomach with formalin.

The gastric mucosa may be damaged by various chemical bacterial and physical factors nervous stimuli and interference with blood supply. The nature of the noxious stimulus its severity duration and portal of entry into the stomach modify the changes occurring in the mucosa. Maintenance of the normal gastric epithelial lining is a dynamic process and simple increase in the rate of regeneration of the cells may cope with increased loss of cells due to some damaging agent without actual ulceration but a point is reached when necrosis and desquamation occur too rapidly for immediate regeneration to occur and erosions or more severe ulceration appear.

The changes characteristically involve the mucosa diffusely with areas of pronounced hyperaemia rendered patchy by mucopurulent exudate mucosal oedema and submucosal petechiae. When damage is severe erosions ulceration and even corrosion of the gastric wall may occur.

The *microscopic changes* in acute gastritis have been described in

biopsy material by E. Palmer (1951) and Motteram (1951). Increased regeneration of cells is reflected by the presence of irregularity of the cell size, nuclear size and staining, poor cytoplasmic differentiation and arrangement of the superficial epithelial cells. These cells may be stratified with indistinct cell borders and containing little or no secretory material in their cytoplasm. The changes are often clearly seen occurring in the depths of the pits. There is oedema of the superficial *lamina propria* which contains increased numbers of plasma cells and lymphocytes and many polymorphonuclear cells (polymorphs) which penetrate the epithelium. Haemorrhages may occur into the interstitium. The overall appearance in some cases may resemble chronic superficial gastritis.




Fig. 7. *Acute gastritis*. Gastric biopsy from a male patient aged 57 years 3 days after severe alcoholic bout followed by haematemesis. Necrotic superficial epithelial cells are visible in the surface mucus. They are regular in height and show poor cell outline. In many the mucous theca is absent. The *lamina propria* is oedematous and infiltrated with plasma cells, lymphocytes and polymorphs (oil and haematoxylin  $\times 600$ ).

More intense stimuli result in frank necrosis of cells which become desquamated into the lumen of the pits and the lining mucus along with migrating inflammatory cells (fig. 7). These necrotic foci may desquamate to leave small erosions or even

penetrate the *muscularis mucosae* to become a frank ulcer. Williams (1950) considered that erosions were initially produced by the rupture of mucosal capillaries through the superficial epithelium. The most severe erosive processes penetrate the gastric wall following necrosis of all coats.

When damage to the stomach has been confined to the superficial epithelium of the ridges and pits surface regeneration is probably always complete (E. Palmer 1951). However it is unlikely that regeneration of damaged glandular cells can occur to the same extent and these cells may permanently disappear or be replaced by non specific or mucus secreting epithelium termed pseudopyloric glands. Damage to the submucosa muscle and serous coats may be followed by the formation of granulation tissue and even deformity of the organ particularly in the antral region.

### Symptomatology

Acute gastritis of minor severity is frequently encountered in clinical medicine the symptoms lasting from a few hours to 2 or 3 days. Anorexia, nausea, occasional vomiting and diffuse epigastric discomfort and tenderness are the outstanding features.

In a severe case the onset is usually sudden with pronounced epigastric pain, persistent nausea and vomiting perhaps with traces of blood. On occasions there may be a profuse haematemesis. The patient is weak, thirsty, drowsy and soon shows the effects of toxæmia, dehydration and electrolyte depletion. He is pale with sunken eyes, inelastic skin, rapid pulse, low blood pressure, muscular cramps and excessive thirst. The urine is scanty, deficient in chlorides and may contain a trace of albumen. The temperature may be subnormal at first and later elevated. There may be constipation or diarrhoea. The abdomen is usually sunken but later may become distended. The tenderness is diffuse with maximum intensity in the epigastrium. The nature and duration of the symptoms are largely determined by the cause of the acute gastritis. Thus in acute staphylococcal food poisoning due to the ingestion of staphylococcal enterotoxin the gastric symptoms predominate whereas in acute staphylococcal gastroenteritis and in *Salmonella* infections the intestinal symptoms predominate with diarrhoea and generalised abdominal tenderness.

E. Palmer (1951) made a valuable study of the stomach in

acute exogenous (staphylococcal) gastroenteritis due to ingestion of food heavily contaminated by staphylococci and their toxins. He investigated 42 previously healthy male subjects aged between 17 and 47 years 38 of whom had been involved in 1 of 3 major outbreaks of 'food poisoning'. On most of them a gastric biopsy was performed with a modified flexible gastric biopsy tube and followed with gastroscopy. The subjects were examined in this way at intervals varying between 5 and 580 hours after the onset of the symptoms. These symptoms usually consisted of vomiting retching chills fever and perhaps some circulatory collapse. Six patients were examined in the acute stage 5-12 hours after the onset and all showed abnormal gastroscopic appearances the majority having hyperaemia oedema erosions petechiae and purulent exudate. The same 6 patients were subjected to gastric biopsy during the acute stage and 5 showed pronounced histological changes the remaining patient was examined 5 hours after the onset of symptoms and showed no histological change but at gastroscopy there was congestion of the mucosa.

The essential histological lesion in these 5 early biopsies was congestion and oedema followed by necrosis of cells at the junction of the pits and glands the damaged cells then being extruded to the surface with exfoliation and erosion. Regeneration of the epithelium was accompanied by interstitial inflammatory cell infiltration.

Twenty four patients were subjected to gastric biopsy and an abnormal gastric mucosa was present in 5 of 6 examined in the first 12 hours all of 4 examined between 26-40 hours 4 of 7 examined between 50-82 hours and 0 of 7 examined between 92-220 hours.

These splendid studies of E. Palmer would therefore suggest that in relatively young individuals a single severe attack of acute gastritis causes considerable inflammatory changes in the gastric mucosa and this is followed by complete healing. It may be that if the attacks were frequent if the subjects were older or if greater numbers had been studied then evidence of the persisting changes of chronic gastritis would be forthcoming. No studies of gastric secretion were reported in E. Palmer's series but it is anticipated that there would have been a depression of acid and pepsin secretion during the acute phase with a return to normal levels with the subsidence of the inflammatory changes.

E Palmer (1934) has also studied the acute gastric symptoms and the gastroscopic and biopsy findings in cases of acute alcoholism (see Chapter 3). He concluded that following an acute alcoholic bout there was evidence of acute gastritis, but that usually there was complete resolution.

However it is our experience that if the alcoholic bouts are severe and often repeated then persistent change may be established the mucosa changing from acute superficial gastritis to chronic atrophic gastritis and perhaps after the course of many years even to gastric atrophy. The factor of chronic malnutrition also plays a part (Joske and Turner 1952) (*cide infra*).

## Aetiology

A number of chemical dietary bacterial as well as psychic stresses may produce acute gastritis. The corrosive action of agents such as caustic soda phenol and mineral acids gives rise to an acute corrosive gastritis in which the stomach wall may perforate. Should the patient survive healing with fibrosis and distortion of the stomach occurs.

In Australia the commonest dietary irritants implicated in acute gastritis are the strong alcoholic beverages. Motteram (1951) illustrated the histological features of the gastric mucosa of a patient from whom a gastric biopsy was obtained 3 days after a haematemesis brought on by a severe bout of alcoholism. Necrotic surface epithelium was present with replacement by actively proliferating epithelial cells growing out from the pits while red cell extravasations were visible in the *lamina propria* (fig. 7).

Finckh *et al* (1952) examined the gastric biopsies of 39 chronic alcoholics. Abnormal gastric mucosa was found in 18. Of these superficial gastritis was present in 14 and atrophic gastritis in 4. Of the cases with superficial gastritis 5 showed subsequent resolution suggesting that the lesion had been acute and reversible.

Later similar figures were obtained in our unit by Joske *et al* (1955) who found 51 out of 95 gastric biopsies obtained from alcoholic subjects showed evidence of gastritis. Williams (1956) found similar changes in a group of 25 alcoholics submitted to gastric biopsy. He also carried out animal experiments which suggested that alcohol consumed in the concentrations found in fortified wines and spirits may cause haemorrhage erosions and ulceration especially if taken on an empty stomach.

Nyfeldt and Vimtrup (1932) examined the stomach of children dying of diphtheria and found evidence of acute gastritis. This was considered to be due to circulating diphtheria toxin for when the stomachs were fixed soon after death and subsequently examined there was no evidence of diphtheritic bacterial invasion. Thomsen (1925) produced gastritis in dogs by the injection of diphtheria toxin. Vimtrup (1929) also found acute gastritis at the post mortem examinations of patients dying from influenza and pneumonia.

As previously stated E. Palmer (1951) performed gastroscopy and gastric biopsy on 42 patients with acute staphylococcal food poisoning and observed the gastritis beginning some hours after the onset of symptoms and usually disappearing within 2 to 4 days without residual damage to the mucosa.

### Treatment

The treatment of acute gastritis depends on the cause and severity of the attack. In the commonest form which is associated with mild food poisoning rest, free fluids and probably a saline purge is all that is required.

The severe attacks may be associated with a specific bowel infection and the causal organism should be sought by culture of a fresh specimen of stool. Transfer to hospital is essential. Antibiotics and sedatives together with the intravenous infusion of electrolytes, glucose and perhaps blood or serum may be indicated. As the patient improves increasing oral feeding is given.

## CHAPTER 3

# CHRONIC GASTRITIS

### Structural changes

Reports of chronic inflammatory changes in the gastric mucosa were criticised and discredited for many years on the grounds that they were caused by post mortem autodigestion. Satisfactory post mortem material was first obtained by Faber and Bloch (1900) using intraperitoneal injection of formalin immediately after death. The pathology of chronic gastritis has since been amply demonstrated by Faber (1935).

The results of the examination of specimens of stomach obtained at operation were recorded by Konjetzny (1928) but have been criticised by Schindler (1947). Schindler *et al* (1939) and Sanders and Mecray (1941) showed that acute erosions of the mucosa could develop during the actual resection of the specimen. Schindler and Ortmyer (1942) obtained stomach biopsies through an abdominal incision but of course this procedure is unsuitable for routine use.

Chronic inflammatory lesions were described in biopsy fragments obtained through oesophageal tubes by Swalm *et al* (1936) while Kenamore *et al* (1946), Benedict (1948) and d Almeida (1948) recorded similar findings in mucosal biopsies. The use of a flexible gastric biopsy tube (Wood *et al* 1949A) has enabled serial biopsies to be made on patients with chronic gastritis immediate fixation of the fragments ensuring freedom from autolysis.

In chronic gastritis there is a wide variety and a continuous spectrum of structural changes. Attempts to classify these changes into different types tend to overlook the variations of distribution, severity and activity which may occur in the same patient over a period of time. A relationship to the clinical features of gastritis is best achieved by classifying according to alterations in the thickness of the mucosa, the concentration of chief and parietal cells and the extent and activity of the inflammatory processes. Thus three main types of gastritis can be recognised.

1 *Superficial gastritis* Here there is superficial inflammation of the mucosa without atrophy of the glands. The mucosa is of normal width and while the inflammatory changes may be slight moderate or severe the lesion is capable of complete resolution. However the changes may persist as a chronic superficial gastritis or with recurrences may progress to atrophic gastritis.

2 *Atrophic gastritis* There is a more diffuse inflammation of the mucosa showing varying degrees of activity coupled with slight moderate or severe atrophy of the specific parietal and chief cells usually leading to a decrease in the width of the mucosa. While the inflammatory changes may resolve to some degree there are permanent changes in the epithelial structures and only partial regeneration of the parietal and chief cells occurs. Recurrent exacerbations of inflammation lead to a severe atrophy which may closely resemble primary gastric atrophy (see Chapter 5).

3 *Gastric atrophy* This is the classical change seen in pernicious anaemia and subacute combined degeneration of the cord. It is included here for the sake of completeness as this change may result from the subsidence of the inflammatory changes in a severe case of atrophic gastritis or alternatively it may be due to a primary atrophy of the mucosa. This is discussed later together with the histological picture (see Chapter 5).

*Diffuse giant hypertrophic gastritis* This is a further rare variety which is described in Chapter 4. It probably represents a different disease to chronic gastritis.

*Macroscopic appearance in chronic gastritis* There may be little alteration from normal appearance. In severe cases the mucosal surface may appear granular or mammillated the mucosa thin and the rugae small (fig. 8). The granularity may be the result of oedema cellular infiltration lymphoid infiltration or cystic change in the epithelial structures. However the degree of distension of the stomach must always be considered when estimating mucosal thickness and the size of rugae—lack of appreciation of this has led surgeons gastroscopists and radiologists to make an erroneous diagnosis of hypertrophic gastritis (fig. 9).

Hebbel (1949) found wide variation in the distribution of chronic gastritis. Local changes were common in the antrum and along the lesser curvature and the presence of changes in these sites was not necessarily indicative of more diffuse changes in the mucosa. However he claimed that changes along the greater curvature






Fig 8 *Chronic atrophic gastritis* Stomach obtained *post mortem* from a male patient aged 64 years with severe chronic atrophic gastritis previously proven by gastric biopsy. The mucosa is smooth, finely granular or slightly nodular. The rugae are flattened and have largely disappeared, an appearance which may have been caused by terminal gastric dilatation (natural size).



Fig 9 *Normal gastric mucosa* in contracted stomach fixed soon after death. The rugae are full and regular. With distension of the stomach they become less prominent (natural size).

occurred usually as part of a more diffuse process in the body of the stomach

*Microscopic appearances in chronic gastritis* : The changes in the gastric biopsy may range from slight superficial increase in plasma cell infiltration of the *lamina propria* to gross alteration in the epithelial structure disappearance of the specific glands and infiltration and replacement by mesenchymal elements. These appearances are best considered under the headings of superficial gastritis and atrophic gastritis although there is no clear dividing line between the two

*Superficial gastritis* The most constant feature is the presence of inflammatory cells in the superficial regions of the *lamina propria*. Where there is only a slight increase of plasma cells unaccompanied by superficial epithelial changes this may be considered to be within physiological limits. The pathological range is considered to be reached when there is more pronounced plasma cell infiltration polymorphonuclear cells are seen migrating singly or in groups from the *lamina propria* to penetrate the superficial epithelium and lie in the pits or superficial mucus (fig 10). Some red cells may be extravasated into the superficial *lamina propria* and there is lymphocytic infiltration of the deeper layer (fig 11).

In acute exacerbations of the gastritis superficial epithelial cell damage is sometimes evident as groups of poorly staining cells in the adherent mucus (fig 7) but a more common feature is the evidence of cell regeneration in the form of irregular or smaller flattened cells. Sometimes these epithelial cells are stratified and show indistinct cell borders poor differentiation and little or no secretory material in their cytoplasm the nuclei vary in size often being large and hyperchromatic.

These changes are sometimes most pronounced in the pits where there is dilatation or cystic change lengthening tortuosity and distortion due to extension of the regenerating cells down into the glands as well as up to the surface. Sometimes this is due to obstruction of the lumen of the pit. The basal two thirds of the mucosa are normal and the total width of the mucosa is unchanged. Should it be possible to remove the damaging agent the mucosa may return to a normal appearance in 2 to 3 weeks.

The histological appearance of chronic superficial gastritis may be identical with that of acute gastritis. Chronicity of the lesion




Fig 10 Active superficial gastritis showing intensive polymorphonuclear infiltration of the lamina propria with penetration of the superficial epithelium by the polymorphs (eosin and haematoxylin  $\times 400$ )



Fig 11 Gastric biopsy showing superficial gastritis. The superficial epithelium is flattened and irregular. There is infiltration of the superficial lamina propria by plasma cells and polymorphs. There are some lymphocytes deeper in the lamina propria (eosin and haematoxylin  $\times 150$ )

is only established by the duration of the symptoms or more convincingly by repeated biopsy when persistent superficial gastritis or perhaps progression to atrophic gastritis is seen

*Atrophic gastritis* In atrophic gastritis the changes in the superficial portion of the mucosa are identical with those in superficial gastritis but the inflammatory changes now extend to involve the lower two thirds of the mucosa with slight (fig 12) moderate (fig 5) or severe atrophy of the glands (figs 13 14 15) The average depth of the mucosa in the processed section varies from 0.47 mm in gastritis with slight atrophy to 0.45 mm in those with severe atrophy (Jo ke *et al* 1935)

Groups of plasma cells and polymorphs are seen in the superficial *lamina propria* while lymphocytes are more prevalent in the deeper areas where they may be aggregated into follicles near the base of the mucosa Occasionally fat cells develop close to these follicles (fig 16) The intervening areas of *lamina propria* contain a delicate connective tissue network which may be oedematous but dense collagenous fibrosis does not occur Sometimes these mesenchymal elements replace the glands completely and often separate the fibres of the *muscularis mucosae* which becomes thickened (fig 17)

The irregularities of the superficial epithelium extend into the glands and the chief and parietal cells are partially or wholly replaced by mucus secreting or non specific cells arranged in coiled pseudopyloric glands a form of post inflammatory metaplasia of the epithelium (fig 18)

Another form of metaplasia is to an intestinal type of gland (Magnus 1937) which is characterised by the appearance of a superficial epithelium containing tall columnar cells with a darkly staining surface interspaced with goblet cells (fig 18) They are arranged to form a crypt with argentaffin cells and at the base Paneth cells The Paneth cells contain coarse eosinophilic refractile granules which stain a dark red with the trichrome stain (fig 19) By means of silver stains argentaffin cells can be demonstrated lying laterally in these crypts (fig 20) (Magnus 1937) This metaplasia is patchy in character more confined to the pits and usually seen to occur abruptly in a single high power field (fig 21)

In spite of these atrophic changes in the glands the mucosa may not be decreased in depth owing to infiltration with inflammatory cells and oedema




Fig. 12 Gastric biopsy showing chronic gastritis with slight atrophy of the glands (trichrome stain  $\times 150$ )



Fig. 13 Chronic atrophic gastritis with severe atrophy. A female aged 64 years with recurrent abdominal pain, anaemia and achlorhydria for 22 years following an episode of gross malnutrition. Gastric biopsy showing severe inflammatory infiltration of the lamina propria with polymorphous plasma cells and lymphocytes. A lymph follicle has formed at the right. The specific glands have almost completely disappeared. During the ensuing 8 years serial biopsies and test meals remained unchanged (eosin and haematoxylin  $\times 150$ )



Fig 14 *Chronic atrophic gastritis*. Gastric biopsy from same case as fig 13. The biopsy shows distortion and elongation of the pits, disappearance of most of the glandular tubules and a severe inflammatory infiltration of the lamina propria (trichrome stain  $\times 150$ ).




Fig 15 *Chronic atrophic gastritis* showing cystic change in the pits with pseudopyloric metaplasia of the glands (eosin and haematoxylin  $\times 150$ ).



Fig. 16 Chronic atrophic gastritis. Lymphoid tissue and fat are replacing the specific glands (eosin and haematoxylin  $\times 180$ )




Fig. 17 Chronic gastritis. There is splintering and thickening of the muscularis mucosae by inflammatory cells (eosin and haematoxylin  $\times 180$ )

With gross atrophy of the glands and post inflammatory metaplasia the end result resembles primary gastric atrophy. Differentiation between atrophic gastritis and gastric atrophy is made on the histological characteristics (Table 3 Chapter 5)

### Symptomatology of chronic atrophic gastritis

The symptoms of chronic atrophic gastritis are distinctive and show but little variation from patient to patient. The patient is usually a female over 40 years of age. Doig and Wood (1952) found a female male ratio of 5:4 and this female predominance was increased when the alcoholics were excluded. The patient's discomfort varies from a sense of excessive fullness in the epigastrium especially after taking food to a dull burning pain. The site of her distress is diffusely situated in the epigastrium. In contrast to peptic ulcer where the patient usually demonstrates the site of the pain and tenderness by pressing in deeply with the tips of the fingers she rubs the epigastrium from side to side with the flat of her hand saying 'This is where I have the discomfort (or the pain) across here'. Rarely pain may also be felt in the back and thus may mimic a penetrating ulcer, chronic pancreatitis or cancer. Like gastric ulcer the pain of gastritis most frequently occurs immediately after taking food or within half an hour.

It is paradoxical that although most of the patients have hypochlorhydria or achlorhydria their discomfort is relieved by taking alkaline powder. Hydrochloric acid taken with their meals may produce relief but in our experience most patients claim they receive no benefit or are made worse—they usually prefer alkaline powder.

Certain foods may aggravate the pain particularly fat, pineapple, cabbage and condiments although there is great variation from patient to patient. There are exacerbations and remissions in the symptoms the exacerbations being brought on by an associated illness, undue mental or physical stress or over indulgence in alcohol. Lack of appetite is common and may lead to loss of strength and weight, this in turn may create a fear of cancer in the mind of the patient and indeed in the mind of the physician. Adding to this fear may be the finding of hypochromic anaemia and occult blood in the stools resulting from gastric erosions.

Occasionally especially in alcoholic gastritis haematemesis or melaena is encountered and thus gastritis must be considered a





Fig 18 Types of mucus secreting cell in chronic gastritis from same biopsy as fig 5. On the upper left hand side the normal mucigenic theca of the superficial gastric epithelium is visible. On the upper right hand side the goblet cells in areas of intestinal metaplasia stain heavily. In the base of the mucosa pseudopyloric mucus secreting cells are increased in amount replacing parietal and chief cells (mucicarmine  $\times 150$ )




Fig 19 Chronic atrophic gastritis. Intestinal metaplasia. A crypt containing dark intestinal type tall columnar epithelium and some goblet cells. Dark granules are visible in the Paneth cells (trichrome stain  $\times 600$ )




Fig. 20. Chronic atrophic gastritis. Intestinal metaplasia. Argentaffin cells are visible in the base of the columnar cells wedged in between them (Masson's silver impregnation  $\times 600$ ).



Fig. 21. Chronic atrophic gastritis. Intestinal metaplasia. Gastric biopsy with a focus of intestinal metaplasia in the superficial epithelium. There is an abrupt change from the normal regular tall columnar gastric epithelium on the right hand side to the deeper staining columnar cells with the dark border and goblet cells on the left hand side (eosin and haematoxylin  $\times 600$ ).



Fig 18 Types of mucus-secreting cell in chronic gastritis from same biopsy as fig 5. On the upper left hand side the normal mucigenic theca of the superficial gastric epithelium is visible. On the upper right hand side the goblet cells in areas of intestinal metaplasia stain heavily. In the base of the mucosa pseudopyloric mucus-secreting cells are increased in amount replacing parietal and chief cells (mucicarmine  $\times 150$ ).



Fig 19 Chronic atrophic gastritis. Intestinal metaplasia. A crypt containing dark intestinal-type tall columnar epithelium and some goblet cells. Dark granules are visible in the Paneth cells (trichrome stain  $\times 600$ ).

aged. She is intelligent but may wear a worried expression. There may be cracks at the angle of the mouth and atrophy of the tongue. The finger nails are brittle with longitudinal furrows rarely koilonychia. There is diffuse epigastric tenderness but no mass or rigidity. The liver and spleen are not palpable except when pronounced malnutrition is present usually due to chronic alcoholism. Examination of the nervous system reveals no abnormality except perhaps some diminution of the knee and ankle jerks.

Although the above description does represent a proportion of cases many reveal no abnormal findings on physical examination except perhaps slight epigastric tenderness during an exacerbation in the disease.

### Aetiology of chronic atrophic gastritis

Chronic atrophic gastritis is a more advanced stage than superficial gastritis and results from some damaging processes either intermittent or continuous the nature of which is often obscure. Once the gastric mucosa has reached the chronic atrophic stage usually there are few signs of recovery of the mucosa as revealed by improvement in the test meal or the gastric biopsy.

Factors which may play a part in the production of chronic atrophic gastritis are sex advancing age perhaps due to increasing atherosclerosis with depleted blood supply of the gastric mucosa chronic debilitating diseases chronic alcoholism associated with malnutrition and the drinking of other gastric irritants prolonged psychic stress and finally irradiation of the stomach especially X ray irradiation either in the treatment of abdominal malignancy or in the purposeful production of achlorhydria in the treatment of duodenal ulcer.

It is of interest to consider some of these factors in greater detail.

TABLE I *Sex distribution in chronic gastritis diagnosed by gastric biopsy*  
There was a preponderance of females especially amongst the cases of superficial gastritis (Dong and Wood 1952)

	Males	Females	% Females
Superficial gastritis	12	36	75
Atrophic gastritis	28	36	56
Total	40	72	64

cause of overt upper gastrointestinal bleeding. However it is most unusual for the bleeding to be of sufficient severity to warrant a laparotomy. Severe and persistent bleeding favours the diagnosis of a peptic ulcer, cancer or bleeding oesophageal varices.

The bowels function normally or there may be intermittent constipation. Although hypochlorhydria or achlorhydria is the rule, attacks of diarrhoea are rare.

The age of onset of the pain is of interest. Doig and Wood (1952) found that whereas the pain of duodenal ulcer usually began in the mid thirties the pain of gastric ulcer, superficial gastritis and atrophic gastritis usually began in the mid forties. When the age of first examination in hospital was studied it was found that patients with superficial gastritis were younger than those with atrophic gastritis. Doig and Wood concluded that gastric ulcer is more closely related to gastritis than to duodenal ulcer and that atrophic gastritis is a more advanced stage of superficial gastritis.

In a survey of 623 patients in the Royal Melbourne Hospital mostly gathered from our unit which was primarily interested in gastroenterology Joske, Finckh and Wood (1953) found 221 cases with pronounced atrophic gastritis. In 40 per cent of these there were gastro intestinal symptoms considered to be entirely or in part due to the gastritis whereas 23 per cent had gastro intestinal symptoms due to other causes and 37 per cent had no gastro intestinal symptoms.

In 50 patients the symptoms were considered to be entirely due to the gastritis. It is our firm belief based on extensive experience with proven cases of chronic atrophic gastritis that these subjects may suffer pain and flatulence due to the inflammatory reaction in the gastric mucosa and to deny this cause of dyspepsia is a grave error of medical judgement. It is obvious from the above survey that chronic atrophic gastritis may be present without symptoms but of course this in no way invalidates the belief that pain may occur in other subjects and that occasionally it may be moderately severe. It should be recalled that gall stones, chronic pancreatitis and even pronounced peptic ulceration may be present without symptoms and it could never be claimed that these diseases rarely or never produce discomfort or pain.

*Physical examination* Although chronic atrophic gastritis occurs in both sexes and even in the third decade it is typically seen in an elderly female with grey hair and perhaps prematurely

gastritis. These may be explained partly on the basis of malnutrition partly by circulating toxins damaging the mucosa. In Sjogrens disease where there is atrophy of the lacrimal and salivary glands with depleted secretion and recurring inflammation in the eyes and mouth gastric biopsy may reveal gastritis with varying degrees of atrophy (fig 22) (Joske *et al* 1955).

It is of interest that in haemochromatosis there is deposition of iron in the deeper layers of the gland cells (Althausen Doig *et al* 1951) but as a rule there are few degenerative changes (fig 23). In 15 gastric biopsies Joske *et al* (1955) found iron deposits in all yet 9 were otherwise within normal limits and 6 showed only superficial gastritis. This is in contrast to the changes in the liver and pancreas in this disease.

*Chronic gastritis due to chronic alcoholism and malnutrition*  
Chronic alcoholism is quite frequently associated with varying degrees of gastritis although some individuals can tolerate excessive intake of alcohol without developing persistent changes. It is probable that after an alcoholic bout the mucosa shows the signs of acute gastritis but it heals without residual change. Thus E. Palmer (1954) selected 34 young men who were physically fit and not alcoholics and examined them by gastroscopy and gastric biopsy within 6 hours of an alcoholic bout. Thirty showed acute exogenous gastritis to gastroscopy and all 34 had abnormal biopsy findings mostly superficial gastritis and resembling the acute changes seen in acute staphylococcal food poisoning (E. Palmer 1951). Eleven of the 34 were submitted to a second biopsy after an interval of 7-20 days during which time they took no alcohol. Nine of the biopsies were now within normal limits.

Palmer (1954) also studied the history of alcoholism amongst 200 patients selected at random whose gastric biopsy was then within normal limits and found that during the previous 12 months 112 admitted being drunk at least once, 81 at least 12 times and 18 at least once a week. Palmer wisely accepted their statements with caution. However it would seem that excess consumption of alcohol does not always cause chronic gastritis. This of course does not indicate that it may not do so in certain susceptible individuals especially when malnutrition is a pronounced feature. Palmer does not relate whether there were other features of chronic alcoholism in these subjects such as disease of the brain, peripheral nerves or liver. However he concludes by favouring

*Sex* Sex appears to play a role in the aetiology of gastritis for females are more commonly affected than males even though alcoholism favours the male incidence 71 per cent of the alcoholics in our unit being males. Doig and Wood (1952) reported that amongst cases of superficial and atrophic gastritis 64 per cent were females (Table 1). This increased incidence amongst females suggests an endocrine factor or perhaps malnutrition as Australian housewives are prone to favour a diet high in carbohydrate and deficient in protein and vitamins (Epstein *et al* 1950). However Fairley, Joske and Turner (1955) could not find convincing evidence that malnutrition played a leading role.

*Ageing and the gastric mucosa* The age of onset of the symptoms of gastritis varies greatly the average being 44 years (Doig and Wood 1952). Our experience indicates that atrophic gastritis results from a progression from superficial gastritis and that once the stage of atrophic gastritis is reached the condition rarely reverses with healing. It is therefore to be expected that gastritis becomes more frequent with advancing age. That it is by no means invariably present in elderly subjects has been revealed by L. Palmer (1954) who carried out gastric biopsy in subjects over the age of 60 who were not suffering any dyspepsia and found 30 with normal gastric mucosa.

Moreover Joske and his colleagues (1955) performed gastric biopsy on 214 patients over the age of 60 and found 18 per cent had normal gastric mucosa. Many of the patients in this series were examined because chronic gastritis was suspected so there was a bias towards an abnormal gastric biopsy finding. It is of interest that in the same series 218 subjects aged between 20 and 40 years were examined by gastric biopsy and 105 (47 per cent) had normal gastric mucosa. Although most of the subjects in this group were suffering from dyspepsia the group included cases of duodenal ulcer where the biopsy is usually normal and also normal subjects without dyspepsia who acted as controls so there was less bias towards an abnormal mucosa.

When achlorhydria is found in an elderly individual it is probably due to chronic gastritis and it should not be considered to be a normal finding any more than chronic nephritis or chronic bronchitis is considered to be normal in the aged.

*Chronic debilitating diseases* Diseases such as chronic infections, ulcerative colitis, rheumatoid arthritis, chronic nephritis and long standing malignancy may be associated with chronic atrophic

the idea that although acute alcoholism may cause acute gastritis chronic gastritis rarely if ever is a direct result of chronic alcoholism. We disagree most emphatically with his opinion.

Indeed Doig and Wood (1932A) surveyed 112 cases of gastritis and found 19 were chronic alcoholics. This series was later extended by Joske, Finckh and Wood (1955) who reported that of 95 biopsies on alcoholic subjects 39 showed superficial gastritis, 11 atrophic gastritis and 1 gastric atrophy without overt pernicious anaemia. Our experience with serial biopsies over many months indicates that whereas superficial gastritis amongst alcoholics may resolve should the patient quit his alcoholic indulgences, a rare happening, it more often progresses to severe atrophic gastritis with continuing alcoholic bouts. Moreover the severe atrophic gastritis suffered by a chronic alcoholic rarely resolves with regeneration of the chief and parietal secreting cells even though the intake of alcohol be reduced or even withheld for a year or more—the chronic gastritis has passed the point of no return.

Whether alcohol acts directly on the gastric mucosa either by bathing and irritating its surface or by way of the blood stream is a matter of debate and it may well be that it acts by both these means. Moreover chronic alcoholism is frequently associated with chronic malnutrition, deficiencies in protein and vitamins of the B group being most commonly encountered. These deficiencies probably play a part in the atrophic changes in the gastric mucosa just as may be clearly seen in the tongue.

In this regard it is of interest to note that Fairley *et al* (1955) carried out a dietetic survey on 30 cases of chronic atrophic gastritis followed for a period of 5 years or more. Although they could not determine a clear relationship between chronic atrophic gastritis and malnutrition they found that 5 patients had taken a consistently poor diet and 9 a poor diet for several months. It is anticipated that studies on a greater number of cases will provide further evidence that malnutrition may lead to atrophic gastritis and that even though a suitable diet with added vitamins is instituted the change in the gastric mucosa may be irreversible.

*Hot tea and gastritis* Edwards and Edwards (1956) have studied tea drinking amongst 155 subjects with dyspepsia in whom gastric biopsy had recently been performed and they produced evidence that the drinking of excessively hot tea favoured the development of gastritis. On the other hand it may be that the sufferer from chronic gastritis is soothed by hot beverages so that chronic





Fig 22 *Atrophic gastritis in Sjögren's disease* Gastric biopsy from female patient aged 60 years with classical Sjögren's disease including conjunctivitis, dry mouth, arthritis and achlorhydria. There is intense inflammatory infiltration of the *lamina propria*, disappearance of glands and some foci of intestinal metaplasia in the elongated pits (trichrome stain  $\times 150$ )




Fig 23 *Haemochromatosis* Gastric biopsy from a female patient aged 60 years with pigmentation, hepatomegaly and debility. Liver biopsy showed cirrhosis and gross accumulation of iron. The glandular epithelium of the stomach contains iron granules (prussian blue stain  $\times 600$ )




Fig. 24 *Radiation gastritis*. A male patient aged 34 years with a chronic duodenal ulcer was given deep X-ray therapy to the stomach. A gastric biopsy taken 66 days after the therapy shows active gastritis with atrophy. The superficial epithelium is flattened and irregular and shows polymorphonuclear penetration. The pits are elongated and distorted and the glands contain many non-specific or mucous secreting cells. There is a heavy infiltration of the lamina propria by polymorphous plasma cells and lymphocytes. At this stage histamine test meal showed achylia. During the next 2 weeks a serial biopsies and histamine test meal revealed a return to normal (see fig. 25) (eosin and haematoxylin  $\times 150$ ).



Fig. 25 *Radiation gastritis*. (A serial biopsy from same patient as fig. 24 taken 25 weeks after radiation of the stomach when the histamine test meal had returned to normal level). The mucosa is almost within normal limits (eosin and haematoxylin  $\times 150$ ).

gastritis favours the drinking of hot tea rather than the contrary. There is no evidence that tea *per se* is a direct cause of gastritis.

**Psychic stress** It has been shown by Wolf and his colleagues (1948) that psychic stress may cause congestion of the gastric mucosa and also the colonic mucosa and that if the stimulus is severe and prolonged erosions and bleeding may occur. Moreover Gray *et al* (1953) found that stress increased the peptic activity of gastric juice. It is possible therefore that long continued stress may lead to chronic atrophic gastritis. Psychic stress certainly aggravates the symptoms in an established case of gastritis.

**Gastritis due to X ray irradiation** The use of X ray irradiation to reduce the secretion of acid and pepsin in the treatment of duodenal ulcer was first advocated by Bassler in the United States of America in 1909 and later was reported by Holman and Lewis (1941) in Tasmania and Ricketts and his colleagues (1948) in Chicago. In 1957 the Chicago group (Levin *et al* 1957) published a follow up of 723 patients and claimed an improvement in the relapse rate compared with the control series. However there were 11 deaths from cancer in the test series 2-13 years after the irradiation (*vide infra*). In 1948 these workers studied the changes in the gastric mucosa in their duodenal ulcer series following X ray irradiation by taking sections at operation or *post mortem*.

In 1951 serial biopsies with the flexible tube were carried out in similar cases in our unit and the morphological changes following X ray irradiation were correlated with the secretion of acid and pepsin (Doig, Inder and Weiden 1951). One patient with chronic duodenal ulcer had biopsies taken 21, 38, 66, 136, 374 and 465 days following a gastric tissue dose of 1500 r given over a period of 3 weeks (figs 24 and 25). The morphological picture of the gastric mucosa changed from normal to one of atrophic gastritis of moderate severity with corresponding depletion of the acid and pepsin secretion the maximum changes being observed during the first 3 months. There was then a slow return towards normal of the histological changes and secretion with almost complete restoration to normal at the end of 1 year. However the secretions returned to normal levels rather than the high levels which preceded treatment.

This case is representative of the majority treated by us and is in keeping with the findings of the Chicago group. However there is a considerable individual variation some patients proving very susceptible and later showing little or no mucosal regeneration or

when she was working in the X ray department for a period of 8 months. During this time she frequently assisted in prolonged screening examinations to test diaphragmatic movement in severely paralysed patients. There was histamine fast achlorhydria and gastric biopsy revealed severe atrophic gastritis. Her duties were changed and her symptoms gradually improved over the next 3 years but the achlorhydria and atrophic gastritis persisted.

We have had no experience with the radiation hazards other than X rays but it may be that with the coming of the atomic age radiation will play an increasing part as a cause of gastritis.

### Relationship of chronic gastritis to peptic ulcer

Doig and Wood (1952) studied the relationship between the age of onset of gastric ulcer, superficial gastritis and atrophic gastritis and found the average age to be similar being 44, 43 and 46 years respectively whereas the average age of onset of duodenal ulcer was significantly less being 36 years. It may be that gastritis in its superficial and atrophic forms may be related to gastric ulcer but not to duodenal ulcer which may have a different aetiology. If gastritis in its initial stages is patchy and limited in extent it may lead to some superficial ulceration. The remaining normal mucosa may then secrete sufficiently concentrated gastric juice to cause further breakdown of tissue with extension of the ulcer in size and depth. However should the gastritis in the first place be widespread it reduces acid and pepsin secretion and thus minimises the tendency to chronic ulceration.

A gastric ulcer with its zonal gastritis (fig. 27) forms an area of impaired gastric secretion and this tends to limit gastric secretion. Thus in gastric ulcer there is reduced secretion of acid and pepsin compared with duodenal ulcer where the mucosa of the body of the stomach is normal or near normal (Perry *et al.* 1956).

In an investigation of 95 cases of duodenal ulcer Doig and Wood (1952C) found 60 per cent with normal gastric mucosa and 30 per cent with superficial gastritis without atrophy. Cox (1952) studied the secreting bulk of gastric mucosa in patients with duodenal ulcer and found it to be in excess of normal. This has not been apparent in our biopsy studies. Magnus (1952) found the mucosa of the body of the stomach to be normal in most cases of duodenal ulcer but that a slight to moderate degree of antral gastritis might occur.

return of secretions others showing only a minor reaction with rapid return to normal of both structure and secretions With increasing dosage of X rays the changes became more pronounced and more prolonged (Levin *et al* 1957 Brown and Wood 1955)

It is of interest that most patients subjected to radiation gastritis suffer little or no discomfort true pain rarely being reported It may be that the pain nerve endings in the stomach wall have been rendered insensitive by the X ray irradiation

Other studies were made in our unit of cases of duodenal ulcer treated by antroduodenectomy followed by X ray irradiation In antroduodenectomy the antrum and most of the first part of the duodenum including the ulcer were removed followed by end to end anastomosis the irradiation served to reduce the high acid secretion which persisted after the operation (Brown *et al* 1952 Brown and Wood 1955 and 1956) Fig 26 shows that there was pronounced lowering of the acid 3 months after the irradiation as compared with the pre irradiation levels




Fig 26 The effect of X ray irradiation on gastric secretion 2000 r gastric tissue doses were given over a period of 3 weeks to lower the gastric secretion in cases of duodenal ulcer previously treated by antroduodenectomy with resection of the ulcer It will be noted that after an initial depression some cases showed a return of acid secretion

One year after the irradiation and later higher levels were reached in many of the cases while a few remained achlorhydric Serial gastric biopsies on a limited number of cases showed there was a corresponding change in the histology of the gastric mucosa (Brown G *et al* 1952 Brown G and Wood I J 1955)

The following case history illustrates the possibility of gastritis being caused by X ray irradiation A nurse aged 26 years developed considerable flatulent dyspepsia and occasional vomiting

and most probably has preceded the cancer (fig 28). However it may well be that the cancer came first and this caused an extensive zonal gastritis. Possibly this was made worse by the malnutrition which results from the anorexia and dyspepsia of cancer. Joske *et al* (1955) found normal mucosa in only 4 of 20 cases of cancer of the stomach examined by random biopsy.



Fig 28 *Epithelial metaplasia in chronic gastritis*. Gastric biopsy from a male patient aged 61 years known to have chronic atrophic gastritis for 10 years. There is an unusual type of metaplasia of the epithelium with the down growth of irregular tubules lined by tall hyperplastic or atrophic epithelium. There is a local inflammatory infiltration. The patient is being closely observed for development of carcinoma (eosin and haematoxylin  $\times 150$ ).

Fairley *et al* (1955) surveyed 32 patients who for 5 years or more had been shown by gastric biopsy and other examinations to have uncomplicated chronic gastritis. During the period of follow up serial tests were made including test meal, gastric biopsy and barium meal. The gastric biopsies showed little overall change during the period of observation, the pronounced atrophic gastritis persisting with some fluctuation in the cellular infiltration. There was a corresponding persistence in the depletion of acid secretion, 14 of 32 showing either a fall or no change in free acid concentration and the remaining 8 a minor rise. No patient in this group showed evidence of gastric cancer either during the period of

It is most improbable that duodenal ulcer would occur in the presence of an extensive gastritis owing to the lowered acid and pepsin secretion which results from the gastritis. However on rare occasions gastritis may develop long after a duodenal ulcer has entered the chronic irreversible stage with penetration into the pancreas. The anomalous position may then be found where a duodenal ulcer is associated with achlorhydria. Probably most duodenal ulcers heal spontaneously after the development of extensive gastritis.

A similar set of circumstances is provided artificially when gastritis is produced by X ray irradiation of the stomach (*vide supra*). This may lead to healing of the duodenal ulcer.



Fig. 27. Zonal gastritis. The edge of a gastric ulcer showing zonal gastritis. The lymph follicles are increased in number and size and the glands have almost completely disappeared. There is dense submucosal fibrosis. Ulceration is evident (lower right) (eosin and haematoxylin  $\times 40$ ).

### Relationship of chronic gastritis to carcinoma

It has been debated over the years whether chronic gastritis favours the development of cancer. It is known that cancer of the stomach especially when situated in the body of the stomach is usually accompanied by hypochlorhydria often achlorhydria. It could therefore be concluded that extensive gastritis is present

reduction in the secretion of intrinsic factor the most significant results being obtained by studying the absorption of  $^{59}\text{Co}$  labelled vitamin  $\text{B}_{12}$  the addition of intrinsic factor returning the absorption to normal. Using our gastric biopsy tube they found atrophic gastritis in 41 of 50 patients with iron deficiency anaemia 2 further cases having gastric atrophy. Davidson and Markson (1955) had similar results.

Witts (1956) concluded that gastritis and achlorhydria are the direct result of iron deficiency.

Joske *et al* (1955) carried out gastric biopsy in 11 patients with hypochromic microcytic anaemia who were not showing overt bleeding and 6 showed atrophic gastritis 3 superficial gastritis and the remaining 2 were within normal limits. They concluded the achlorhydria so frequently seen in this type of anaemia results from inflammatory gastritis.

*Absorption of iron from alimentary canal.* Using foods containing radioactive iron Moore (1955) showed that absorption of iron was normal in patients suffering from iron deficiency anaemia and achlorhydria. It is most probable that the majority of his subjects had atrophic gastritis. Moreover he demonstrated that the addition of liberal volumes of dilute hydrochloric acid did not increase the absorption.

TABLE 2 *Absorption of iron*

Cases of hypochromic anaemia (probably associated with chronic atrophic gastritis and anaemia) can absorb liberal amounts of iron given by mouth. However cases showing the malabsorption syndrome with steatorrhoea show grossly impaired absorption.

Authority	Test dose of ferrous iron given by mouth	Disease state	% Iron absorbed
Dubach <i>et al</i> (1948)	1 mg/kg body weight	Normal subjects Hypochromic anaemia	2-21 4-71
Badenoch and Callender (1954)	40 mg	Hypochromic anaemia Steatorrhoea and hypochromic anaemia	20+ 0-3



observation or at the final assessment. In addition there were 9 cases of atrophic gastritis who died before the 5 year period had passed and none had gastric cancer. Although the number of patients in this study is limited it does not support the opinion that chronic gastritis strongly favours the development of cancer.

If atrophic gastritis is not a precursor of cancer it would suggest that the causal factor of gastritis is not carcinogenic. However in this regard one would not readily care to discard the thought that X rays could act as a carcinogen (Brown and Wood 1956). Indeed Levin and his colleagues (1957) in a follow up of 723 patients treated by irradiation for duodenal ulcer reported 6 to have died from carcinoma of the stomach, duodenum or pancreas 2-13 years after irradiation. They considered that the significance of this is yet to be determined.

Witts (1956) who had studied cases of achlorhydria, atrophic gastritis and hypochromic anaemia for many years could find no reliable data on the liability of women with iron deficiency and achlorhydria to develop carcinoma of the stomach in later life. He concluded 'we know that they are abnormally prone to develop carcinoma of the mouth and hypopharynx and we should keep an open mind about the liability to carcinoma of the stomach'. Fig. 28 is of interest in this regard.

### Relationship of chronic gastritis to anaemia

When anaemia does occur in cases of chronic atrophic gastritis it is usually of the iron deficiency type, being hypochromic and microcytic. Several factors contribute to this: anorexia leads to a depleted intake of food and the food taken is often low in iron and protein content; secondly it has been claimed that there may be impaired absorption of iron due to greatly reduced secretion of acid by the stomach; thirdly there may be repeated gastric haemorrhages due to excessive shedding of gastric epithelium, the haemorrhage usually being minimal but oft repeated; and lastly, with increasing atrophy of the gastric mucosa the supply of intrinsic factor may be reduced, causing impaired absorption of vitamin B<sub>12</sub> and ultimately pernicious anaemia (see Chapter 5).

**Witts' anaemia.** Witts (1956) found that the incidence of achlorhydria is increased at all ages amongst cases of iron deficiency anaemia—under 50 years it was 4 times as common as in a control group. He and his colleagues at Oxford also found a

medical treatment (fig. 29). However the blood loss in gastritis is usually occult although it may lead to pronounced anaemia.

### Treatment

Since most patients with chronic gastritis suffer minor bouts of discomfort they only require simple treatment during the attacks. We found that alkaline powders provided the greatest relief, odd though this may seem in the light of the achlorhydria. It is of course rational to administer dilute hydrochloric acid with meals, but few claim benefit from this and actually most are made more uncomfortable. However there were exceptions to this experience for a few patients obtained considerable benefit from the acid.

Should the relapse be moderately severe and prolonged oral penicillin may be given with benefit—100 000 units 4 hourly for 10 days. There is evidence that penicillin is absorbed through normal gastric mucosa (Hurley and Dazeley, 1957) but it is not known whether this pertains when chronic gastritis is present.

When overt haemorrhage occurs with melaena or haematemesis the patient should be given rest in bed and placed on a haematemesis regime. However rapid and severe blood loss requiring a liberal blood transfusion or gastrectomy rarely occurs.

Should anaemia be present it is usually of the iron deficiency type and responds to iron given by mouth. Ferrous sulphate is a satisfactory salt but should this cause added dyspepsia the other salts such as ferrous lactate or iron and ammonium citrate may be satisfactory. Some patients are intolerant to all oral medication or the anaemia is not relieved owing to intestinal malabsorption (Table 2). In these an intramuscular preparation such as Imferon® (1 ml twice a week) is of value. Hagedorn (1957) has described a short high dosage course of this preparation.

If the anaemia is not relieved by iron then further search should be made for blood loss or other causes of anaemia such as chronic nephritis (Joske *et al.* 1956) or carcinoma of the colon. It should be remembered that the atrophic gastritis may eventually cause progressively increasing vitamin B<sub>12</sub> deficiency from diminishing secretion of intrinsic factor—macrocytic anaemia will then appear. This onset of pernicious anaemia will respond to injections of B<sub>12</sub>.

When chronic flatulent dyspepsia, lack of appetite, loss of weight and anaemia occur in an elderly patient as a result of

\* Bengel Laboratories Ltd, Cheshire, England.

It will be noted in Table 2 that the absorption of iron from the alimentary canal is increased in iron deficiency anaemia unless this deficiency is associated with steatorrhoea when absorption is reduced. In steatorrhoea it is the intestinal barrier which is at fault for iron is readily absorbed and utilised when injected parenterally (Dubach *et al.* 1948; Badenoch and Callender 1954).

It would appear probable that the most common causes of hypochromic anaemia associated with gastritis are deficiency of iron in the diet and loss of iron from chronic bleeding from the gastric mucosa (fig. 29). Moreover it is considered that the iron deficiency follows the gastritis rather than causing it as has been claimed by Witts. Indeed the gastritis is one of the causes of the iron deficiency.



FIG. 29. Chronic atrophic gastritis with haemorrhage. A male patient aged 38 years had periods of epigastric pain over a period of 5 years. Admitted with haematemesis and melaena. No ulcer or cancer found. The gastric biopsy performed 11 days after admission shows haemorrhage and infiltration of the lamina propria. The red = lysate in dark (1 hr. meth. stain  $\times 150$ ).

**Overt bleeding.** In a study of 112 cases of gastritis diagnosed by gastric biopsy, Doig and Wood (1952) described 4 cases of superficial gastritis and 3 of atrophic gastritis who had been admitted to hospital with haematemesis or melaena. No other cause for the bleeding such as ulcer or cancer was found and all recovered with

## CHAPTER 4

# DIFFUSE GIGANTIC HYPERTROPHIC GASTRITIS

Since the earliest description of diffuse giant hypertrophic gastritis by Menetrier (1888) perusal of the literature shows that personal experience of this disease by any one author is restricted to only a few cases and we have encountered only one case to be described later. Nevertheless Fieber (1955) has recorded an excellent analysis of 50 cases reported in the literature. The lesion is traditionally regarded as inflammatory in origin but the aetiology is unknown. It may be that the pathological changes seen in this disease are basically neoplastic, an anatomical variant of gastric adenomatosis and that the inflammatory reaction is a secondary occurrence. This view is favoured by the very prominent folds, the profuse epithelial proliferation and the occasional development of true carcinoma (Matzner *et al.* 1951; Palumbo *et al.* 1951; Texter *et al.* 1953).

The lesion may diffusely involve the whole gastric mucosa or it may be more localised in distribution, particularly in the antrum. The gastric mucosal folds are swollen, tortuous, nodular and even polypoid, ranging from 0.5 to 3.5 cm in height. The surface of the folds may be congested with a mammillary or cobblestone pattern, superficial haemorrhages and erosions. The mucosa is soft and usually mobile but may become more turgid and relatively fixed due to inflammatory induration or malignant change. There may be a tendency for the redundant mucosa to prolapse through the pylorus.

The *microscopic appearance* is basically that of benign hyperplasia of the surface epithelium resulting in extensive elongation, branching and tortuosity of the pits. Cystic changes may become pronounced, presumably the result of obstruction and inflammation in these pits. There is infiltration of the interstitium by inflammatory cells. Normal glands may be visible in the depth of the mucosa.

chronic gastritis malignancy may well be suspected by both patient and physician. Indeed it is essential first to exclude the presence of cancer in such a patient. But once this has been done, the patient will profit greatly by having the disease entity chronic gastritis explained to him (or more frequently her). He should be reassured that his symptoms are typical of this malady and that there is no evidence of cancer. Moreover, he will be comforted by the knowledge that although he may suffer a minor relapse from time to time he will enjoy good health and regain his weight with improved diet and medicinal treatment. He should be seen from time to time to maintain treatment and to ensure accuracy of diagnosis.

\* \* \*

It is our opinion that the acceptance by physicians and surgeons of chronic gastritis as a real disease entity with specific symptoms and signs is essential to the efficient practice of medicine.




Fig 30 Diffuse giant hypertrophic gastritis Operative specimen of stomach. There is pronounced thickening, tortuosity and maturation of the rugae of the body of the stomach. The antrum (left) is normal. The patient was a male aged 47 years with epigastric pain and recurrent haematemesis for 2 years. A pre-operative diagnosis of carcinoma of the stomach was made in error (approximately half natural size).



Fig 31 Diffuse giant hypertrophic gastritis. Microscopic appearance of the gastric mucosa (see fig 30). There is gross proliferation and atrophic change in the pits. The interstitial tissue is heavily infiltrated with inflammatory cells (eosin and haematoxylin  $\times 45$ ).

Fieber (1955) found that diffuse giant hypertrophic gastritis occurred in adults of all ages but particularly in the fourth to sixth decades of life. Males outnumbered females by 3:1. The average duration of symptoms was 2 years and these included in order of frequency diffuse post prandial epigastric pain and tenderness sometimes relieved by food and alkalis weight loss vomiting and bleeding. Test meal findings were not characteristic. The large rugae were detected by gastroscopy and barium meal examination.

The differential diagnosis at this stage is from (1) malignant neoplasms of the stomach notably carcinoma (most frequent misdiagnosis) sarcoma and reticulosis (2) benign neoplasms of the stomach (3) large gastric rugae especially in the deflated stomach (Ricketts *et al* 1947) (4) granulomata of the stomach (5) extrinsic tumours invading the stomach.

Biopsy through the gastroscope cytological studies and blind biopsy with the flexible gastric biopsy tube may be of value in differentiating between these conditions but a reliable diagnosis of diffuse giant hypertrophic gastritis can only be made by the pathologist with access to the full thickness of the stomach wall at any area he may wish to examine. Cases in which there is any suspicion of a resectable malignancy should be submitted to laparotomy.

Stiegmann *et al* (1957) consider the following features suggest that the large rugae are benign a long history response to medical treatment absence of a palpable mass normal or near normal test meal a smooth outline of the filling defect elasticity of the underlying gastric wall smoothness and continuity of the rugae and decrease in the size of the lesion during observation. Such a case they suggest may undergo conservative treatment and observation but generally the resemblance of diffuse giant hypertrophic gastritis to carcinoma leads to laparotomy.

Fieber (1955) states that the response to medical regimes or deep X ray therapy is not good. Where there is suspicion of malignancy and the lesion is diffuse total gastrectomy is necessary but if the lesion is proven to be benign at operation subtotal gastrectomy suffices even though the lesion is diffuse.

*Case history* We are indebted to Dr W. E. King for the following case record. A male patient aged 42 years was admitted to hospital with a history of indigestion for 10 years and during the past year a haematemesis followed by tiredness malaise and epigastric pain. A barium meal examination showed an extensive

## CHAPTER 5

# GASTRIC ATROPHY, PERNICIOUS ANAEMIA AND SUBACUTE COMBINED DEGENERATION OF THE CORD

### Structural changes of gastric atrophy and pernicious anaemia

In 1849 Thomas Addison presented his classical description of the anaemia which for many years bore his name. It was in 1860 however, that Flint drew attention to the gross reduction of gastric secretion in Addisonian anaemia and Cohn and von Mering (1866) noted the absence of hydrochloric acid from the gastric juice. The gastric atrophy accompanying this condition was first noted by Samuel Fenwick (1870) who subsequently confirmed this finding in 3 other cases (Fenwick 1880). Faber (1935) considered the changes in the stomach were inflammatory in origin and that gastric atrophy was the end result of this process but Mignus and Ungley (1938) described a non-inflammatory atrophy of the body of the stomach in 8 patients with pernicious anaemia. Similar observations were also made by Meulengracht (1939) and Cox (1943).

The macroscopic appearances described by these authors are characteristic of gastric atrophy. The mucosa of the body of the stomach is reduced in thickness and the atrophy also includes the muscle coats. By contrast the pyloric mucosa and muscle coat are normal in appearance the demarcation between the two areas being quite sharp (fig. 32).

The histological changes in the mucosa of the body of the stomach as recorded by these authors resemble the gastric atrophy described in the gastric biopsies by Motteram (1951) of our unit. These changes are the virtual disappearance of the parietal and chief cells of the glands with partial replacement by areas showing intestinal metaplasia and non specific secreting cells while inflammatory changes are minimal or absent (fig. 6). In 90 per cent of cases all the specific cells of the glands have disappeared while



filling defect suggestive of carcinoma of the stomach. At gastroscopy thickened folds suggestive of carcinoma were seen projecting into the stomach. Total gastrectomy was performed by Mr J. O. Smith. The post operative recovery was slow and he continued to suffer from dyspepsia. The pathologist Dr J. D. Hicks reported that the stomach measured 23 cm along the lesser curvature the mucosa was pink hypertrophied and heaped up into numbers of large papillary folds the whole of the mucosa having a gross mammillary appearance (fig. 30). On microscopic examination the epithelium of the mucosal folds was in many places normally differentiated but contained numerous plasma cells in the outer portion of the mucosa and many lymphocytes in the basal portion (fig. 31). Also there were cystic glands in the lower portion of the mucosa and in some areas these were well differentiated. Some dilated glands extended into the submucosa and were lined by regular mucus secreting cell.

Motteram 1950 Doig Motteram Robertson and Wood 1950 Motteram 1951 Wood 1951 Finckh and Wood 1953 Badenoch and Richards 1953 Doig 1954 Badenoch 1954) In practice a continuous range of histology is encountered from atrophic gastritis to gastric atrophy so that it is possible that many cases of gastric atrophy are in fact the inactive end result of a severe atrophic gastritis. Conversely cases of gastric atrophy from time to time may develop superimposed inflammatory changes. For the purposes of classifying and analysing our biopsy results we have used the criteria set out in Table 3.

TABLE 3 *The histological changes in atrophic gastritis and gastric atrophy*

Biopsy examination	Atrophic gastritis	Gastric atrophy
Disappearance of gland cells	Usually incomplete	Usually complete
Intestinal metaplasia	Uncommon and is patchy and often confined to the pits	Usually present and tends to be diffuse
Inflammatory changes	Epithelial irregularity cellular infiltration including polymorphs increase in lamina propria thickened and splintered muscularis mucosae all pronounced	Usually mild or absent with only superficial changes

Of 1000 gastric biopsies reviewed in 1955 by Joske, Finckh and Wood, 100 were obtained from patients with either pernicious anaemia or subacute combined degeneration of the cord or both. Gastric atrophy was present in 40 and in the remaining 60 there were varying degrees of chronic gastritis with severe atrophy. All of the 100 patients had histamine fast achlorhydria. These authors also found gastric atrophy in 16 biopsies from patients without overt pernicious anaemia or subacute combined degeneration of the cord. One of these subsequently developed macrocytic anaemia and subacute combined degeneration of the cord which responded to treatment with vitamin B<sub>12</sub> (Robertson, Wood and Joske 1955) (*vide infra*). Three of the other biopsies were obtained from patients who at other times showed atrophic gastritis illustrating both the relationship between these conditions

an occasional chief cell may be found in the others. Parietal cells are rarely seen. Intestinal metaplasia is usually present and diffusely distributed in the mucosa the superficial epithelium consisting of columnar cells with a dark border goblet cells and argentaffin cells while Paneth cells are present in the bases of the crypts. Gastric type epithelium when present stains intensely for mucin. The superficial epithelium is regular and the *muscularis mucosae* is thin. The *lamina propria* is slightly increased and often contains a few plasma cells and lymphocytes the latter occasionally aggregated into small follicles at the base of the mucosa. The average width of the processed mucosa in this condition is 0.42 mm the average normal being 0.54 mm (Joske *et al* 1955)



Fig 32 *Gastric atrophy.* A male aged 58 years with severe pernicious anaemia died immediately after admission to hospital from an emia bronchopneumonia and cardiac failure. There is gross atrophy of the mucosa of the body of the stomach. The antrum and pylorus (left) appear normal (approximately half normal size).

It cannot be stated however that the morphological changes in the stomach found in cases of pernicious anaemia and subacute combined degeneration of the cord are uniform except in so far as most authors now agree that atrophy with or without inflammation of the mucosa of the body of the stomach is invariable and usually severe (Doig 1949 Doig and Wood 1950 Doig and

gastric biopsies and maximum stimulus histamine test meals over a period of 8 years (fig 33). With treatment he made the usual complete clinical recovery and then remained well on maintenance treatment first with liver extract and then liberal doses of vitamin B<sub>12</sub> but gastric atrophy and achlorhydria persisted throughout (Finckh and Wood 1953).

### Symptomatology of gastric atrophy and pernicious anaemia

Much depends on the sequence of events which leads to gastric atrophy. If gastric atrophy is a slow primary atrophy of the mucosa probably genetically determined and unaccompanied by an inflammatory reaction (gastritis) then the gastric symptoms will usually be minimal or absent and the clinical picture will be dominated by varying degrees of vitamin B<sub>12</sub> deficiency with anaemia and subacute combined degeneration of the cord. If however the advent of vitamin B<sub>12</sub> deficiency is preceded by severe chronic gastritis slowly progressing from the stage of superficial gastritis to atrophic gastritis and perhaps thence to gastric atrophy then it might be expected that there would be a long history of intermittent flatulent dyspepsia typical of gastritis.

Analysis of the symptoms before the appearance of overt pernicious anaemia is therefore of considerable interest.

Most cases of pernicious anaemia enjoy excellent health without dyspeptic symptoms once adequate vitamin B<sub>12</sub> therapy has been established. Thus the gastric atrophy *per se* produces few symptoms. On the other hand some patients first presenting with the pernicious anaemia picture complain of sore tongue, poor appetite, mild to moderate flatulent dyspepsia, general malaise and perhaps bouts of diarrhoea. Most of this can be attributed to vitamin B<sub>12</sub> deficiency and not to gastric atrophy and the resulting achlorhydria.

The male patient aged 63 years previously reported by Robertson, Wood and Joske (1955) is of considerable interest in this regard. When seen in February 1949 he stated that he had suffered from flatulent dyspepsia since the age of 15. The pain came on soon after taking food and was relieved by belching or taking alkaline powders. His appetite was poor and he had lost weight. He was referred to the hospital with the provisional diagnosis of gastric carcinoma but this was excluded by X ray and gastroscopy. Investigation showed that he had achlorhydria, gastric atrophy (repeated biopsies), normal bone marrow and

and the difficulty of completely separating them on histological grounds

Following the administration of vitamin B<sub>12</sub> to cases of pernicious anaemia there is no change in the appearance of the gastric biopsy (Doig and Wood 1950 Finckh and Wood 1953) We have not been able to observe a change in the size of the epithelial cells which would be in accord with the findings of Graham and Rheault (1954) who reported large gastric epithelial cells in gastric washings comparable to the megaloblasts in the bone marrow and which returned to normal size with vitamin B<sub>12</sub> therapy Also Massey and Rubin (1954) and Gamble *et al* (1957) noted no significant change in the size of the epithelial cells in gastric washings following vitamin B<sub>12</sub> therapy




Fig 33 *Gastric atrophy*. A male aged 47 years with pernicious anaemia treated with liver extract and then vitamin B<sub>12</sub> for 8 years. The gastric biopsy shows diffuse intestinal metaplasia and complete atrophy of the gland. There is slight infiltration of the lamina propria with plasma cells and lymphocytes. Serial gastric biopsies taken before and during treatment have shown no change in the histological appearance (mucicarmine  $\times 150$ )

In our experience the gastric atrophy and the resulting achlorhydria remain unchanged even after many years of full parental vitamin B<sub>12</sub> therapy. One such patient a male chocolate maker aged 47 years suffering from pernicious anaemia and subacute combined degeneration of the cord has been studied by serial

cord in the absence of any abnormality in the peripheral blood or bone marrow may present a considerable problem in diagnosis (Victor and Lear, 1956) (figs 34-35). It is odd that vitamin B<sub>12</sub> deficiency causes such a variable clinical picture varying from severe anaemia with no overt subacute combined degeneration of the cord to severe combined cord degeneration and no anaemia. In our present state of knowledge we can but believe that there is a different degree of system vulnerability varying from patient to patient. However histamine fast achlorhydria is always present and gastric biopsy reveals a severe degree of atrophy of the gastric mucosa (Doig *et al.* 1950-1).



Fig. 34. Severe atrophic gastritis. A male patient aged 64 years was admitted with weakness in the legs and evidence of posterior and lateral column degeneration in the spinal cord. There was a histamine fast achlorhydria but peripheral blood and bone marrow were normal. The gastric biopsy shows irregularity and shedding of the superficial epithelium, pseudopyloric metaplasia and complete loss of parietal and chief cells of the glands. The lamina propria is infiltrated with inflammatory cells and the muscularis mucosae is thickened and splintered. There was a good response to vitamin B<sub>12</sub> therapy (eosin and haematoxylin  $\times 150$ ).

\*Cobalt labelled vitamin B<sub>12</sub> has been used to detect impaired vitamin B<sub>12</sub> absorption in this syndrome (McIntyre *et al.* 1956). Liversedge *et al.* (1957) found radioactive vitamin B<sub>12</sub> of considerable help in confirming doubtful cases of subacute combined

peripheral blood and only very minor signs suggestive of subacute combined degeneration of the cord. The technique for estimating serum B<sub>12</sub> levels was not then available. He was not given vitamin B<sub>12</sub> treatment as he was enjoying quite good health and he improved with reassurance and a more satisfactory diet. However he was kept under close observation and in September 1954 he developed overt pernicious anaemia with pronounced subacute combined degeneration of the cord and a low level of serum vitamin B<sub>12</sub>. There was a full response to liberal vitamin B<sub>12</sub> and he was well controlled by continuing therapy. When last seen nearly 4 years later he was enjoying excellent health but still suffered from some flatulent dyspepsia. Examination of his blood and nervous system were satisfactory, there was achlorhydria to maximum histamine stimulation covered by antihistamine (Kay 1953) and gastric biopsy showed the same picture of gastric atrophy.

A similar delay in the development of overt pernicious anaemia after depleted vitamin B<sub>12</sub> absorption from intrinsic factor lack may be observed in patients subjected to total gastrectomy. Here the onset of symptoms may be delayed for periods up to 3 years or more after the gastrectomy, the patient meanwhile subsisting on the body stores of vitamin B<sub>12</sub> (Welbourn *et al* 1956, Harvey 1956).

Recently Holmes (1956) has stressed the mental changes which accompany pernicious anaemia and which he attributes not only to cerebral anoxia but also to the effect of vitamin B<sub>12</sub> deficiency on the brain for the symptoms may be present before anaemia and subacute combined degeneration of the cord appear. More over they are relieved by vitamin B<sub>12</sub> therapy.

Wilkinson (1948) reported flatulence or dyspepsia in 59 per cent of cases of pernicious anaemia and the symptoms often antedated the anaemia by many years. Davidson (1957) noted that poor appetite and vague abdominal discomfort were frequent in pernicious anaemia but diarrhoea only occurred in 9 per cent of 250 cases which he reviewed in Edinburgh during 1944-56. He contrasted this finding with Cabot's series of 1200 cases seen before the discovery of liver therapy where the incidence of diarrhoea was 51 per cent.

It may well be that vitamin B<sub>12</sub> deficiency directly impairs gastric and intestinal function and this is remedied by vitamin B<sub>12</sub> therapy.

The development of subacute combined degeneration of the

severe atrophy of the gastric mucosa so the cause of this atrophy must be sought when determining the cause of pernicious anaemia

Since gastric biopsy examinations were first made in our unit in 1948 we have made a number of studies of pernicious anaemia

In 1950 Doig and Wood examined by gastric biopsy 45 cases of established pernicious anaemia or subacute combined degeneration of the cord and more recently Joske Finckh and Wood (1955) have extended the series to 100 cases. All had severe gastric changes with atrophy 40 were classified as gastric atrophy no signs of inflammation being present and 60 had varying degrees of chronic gastritis with severe atrophy. All had histamine fast achlorhydria

In Scotland Markson and Davidson (1956) carried out gastric biopsy in 16 cases of pernicious anaemia and found gastric atrophy in 14 and atrophic gastritis in 2

It is rational to conclude that severe atrophic gastritis may lead to gastric atrophy with subsidence of the inflammatory reaction and that the causes of pernicious anaemia thus include the causes of superficial gastritis which then progresses to atrophic gastritis (see Chapter 4). This progression probably takes a number of years and so far our 8 year study of gastritis has not produced an example of the complete progression through the 3 phases. We have seen some cases of superficial gastritis progressing to atrophic gastritis and some cases of severe atrophic gastritis progressing to gastric atrophy but without the development of overt pernicious anaemia or subacute combined degeneration of the cord. These may appear later with depletion of the vitamin B<sub>12</sub> stores

There is also evidence that gastric atrophy may occur as a primary process a simple abiotrophy of the gastric mucosa without a preceding inflammatory gastritis (Magnus and Ungley 1938 Motteram 1951). It may well be that there is a genetic influence in this primary degeneration. It is therefore of importance that the occurrence of pernicious anaemia in families has been reported by a number of writers. Wilkinson (1949) investigated 1600 cases of pernicious anaemia and found a history of pernicious anaemia in at least one other member of the family in 19 per cent. Mosbeck (1953) investigated 132 patients with pernicious anaemia in whom a relatively complete family history could be obtained. There were 1958 relatives with 51 (2.3 per cent) cases of pernicious anaemia this was 20 times that found in a control series. More over 30 per cent of the families had one or more cases additional to the index case



degeneration of the cord either when there was no accompanying anaemia or when vitamin B<sub>12</sub> had been given before the blood state had been adequately examined




Fig 35 *Subacute combined degeneration of the cord* A female patient aged 64 years developed urinary retention and progressive weakness of the legs over 2 weeks. She had a smooth tongue, wasting of the leg muscles with absent deep reflexes, extensor plantar responses and loss of sensation to light touch, pinprick and vibration over both legs. Gastric biopsy showed gastric atrophy. She died of urinary sepsis 6 weeks later despite injections of liver extract and antibiotics. A section of the lumbar portion of the spinal cord shows extensive demyelination of the posterior and lateral columns (Weigert stain  $\times 20$ )

### Aetiology of gastric atrophy and pernicious anaemia

It is of considerable importance to speculate as to the cause of pernicious anaemia. As will be discussed later it must be accepted that pernicious anaemia is due to impaired vitamin B<sub>12</sub> absorption which in turn is due to deficiency of Castle's intrinsic factor. This intrinsic factor is present in the mucoprotein fraction of gastric juice. Some patients may also secrete intrinsic factor in the duodenum and perhaps the upper ileum but in relatively minimal amounts (Castle *et al* 1929, Grasbeck 1956). In pernicious anaemia the cause of the deficiency of Castle's intrinsic factor is

of 18 patients with severe atrophy of the mucosa had a low total volume of secretion achlorhydria and low or absent pepsin while all of 36 patients with normal or nearly normal mucosa had normal or intermediate type secretions (Wood *et al* 1949B). Similarly the total mucus secretion was reduced in chronic gastritis and more so in gastric atrophy even though the mucus was more concentrated (Weiden 1949).

In a later survey Joske *et al* (1955) recorded 784 gastric biopsies in patients who had also been examined by histamine test meal (see fig 1 Chapter 1). The test meal yielded a concentration of over 20 units of free acid in 76 per cent of 275 biopsies which were within normal limits in only 15 per cent of 172 showing pronounced atrophic gastritis and in none of 41 showing gastric atrophy.

In 1909 Castle and his co workers demonstrated that an intrinsic factor present in normal human gastric juice combined with the food to produce a haemopoietic response when given orally to patients with pernicious anaemia.

The localisation of the intrinsic factor in the pylorus and duodenum of the pig was achieved by Meulengracht (1939) but Fox and Castle (1942) demonstrated that in man this factor was present mainly in the body of the human stomach so that the gastric lesion could be correlated with the absence of intrinsic factor. Goldhamer (1937) had shown that the gastric juice of patients with pernicious anaemia contained minute amounts of intrinsic factor but because of the achylia he suggested that the defect was mainly quantitative.

Intrinsic factor has since been identified as a mucopolypeptide (Williams *et al* 1954) present in the mucoprotein of gastric juice (Latner *et al* 1954). It now seems likely that intrinsic factor combines with extrinsic factor (vitamin B<sub>12</sub> or cyanocobalamin isolated by Smith 1948 and Riches *et al* 1948) and facilitates its absorption in the small intestine (Ross 1950). It is stored in the liver combined to a beta globulin (Pitney *et al* 1955) and then combined to an alpha globulin (Pitney *et al* 1954). It is transported in the plasma to the sites of utilisation. In conjunction with folic acid vitamin B<sub>12</sub> is thought to play a major role in the synthesis of nucleoprotein particularly at the sites of rapid cell proliferation such as the bone marrow. Vitamin B<sub>12</sub> deficiency occurring as a result of deficiency of intrinsic factor is followed by a decrease in the rate of cellular proliferation. The cells instead of dividing

It is also of interest that Videbaek and Mosbech (1954) found gastric carcinoma to be 4 times higher in the relatives of patients with gastric carcinoma than in a control series and that in their series the incidence of pernicious anaemia was about 3 times higher than the normal in the relatives of patients with gastric carcinoma a finding which accords with the treble incidence of gastric carcinoma in patients with pernicious anaemia. They concluded changes in the gastric wall—which presumably may be the result of exogenous as well as hereditary factors—therefore appear to predispose to both gastric carcinoma and pernicious anaemia.

Doig *et al* (1950B) carried out histamine test meals on 134 healthy medical students and found only one with histamine fast achlorhydria. This student a male aged 25 had no signs of pernicious anaemia and the barium meal X ray was negative. Gastric biopsy in 1949 showed an unusual type of atrophy of the acid and pepsin secreting cells and an absence of inflammation. His maternal grandmother had died at the age of 70 from pernicious anaemia. His mother at the age of 44 suffered from flatulent dyspepsia and weakness and a gruel meal showed hypochlorhydria and excess mucus. In 1955 histamine fast achlorhydria was still present and in 1957 when last seen he was well and had required no treatment.

In conclusion it can be stated that there is evidence that a proportion of the cases of pernicious anaemia result from a primary atrophy of the gastric mucosa which may begin early in life and be determined by a genetic factor. The extent of the atrophy may not be sufficient to produce vitamin B<sub>12</sub> deficiency and pernicious anaemia till middle age or later. Secondly the remaining cases of pernicious anaemia probably result from superficial gastritis progressing to severe atrophic gastritis with resulting depletion of Castle's intrinsic factor which in turn leads to depleted vitamin B<sub>12</sub> absorption and vitamin B<sub>12</sub> deficiency. Thus atrophic gastritis may become quiescent with disappearance of the inflammatory reaction in the gastric mucosa and so the gastric biopsy findings, in that of gastric atrophy.

### **The relationship of the gastric changes to pernicious anaemia subacute combined degeneration of the cord and other megaloblastic anaemias**

In a correlation of the gastric secretions obtained by histamine test meal with the findings on gastric biopsy it was found that all

of 18 patients with severe atrophy of the mucosa had a low total volume of secretion achlorhydria and low or absent pepsin while all of 36 patients with normal or nearly normal mucosa had normal or intermediate type secretions (Wood *et al* 1949B). Similarly the total mucus secretion was reduced in chronic gastritis and more so in gastric atrophy even though the mucus was more concentrated (Weiden 1949).

In a later survey Joske *et al* (1955) recorded 784 gastric biopsies in patients who had also been examined by histamine test meal (see fig 1 Chapter 1). The test meal yielded a concentration of over 20 units of free acid in 76 per cent of 275 biopsies which were within normal limits in only 15 per cent of 172 showing pronounced atrophic gastritis and in none of 41 showing gastric atrophy.

In 1929 Castle and his co workers demonstrated that an intrinsic factor present in normal human gastric juice combined with the food to produce a haemopoietic response when given orally to patients with pernicious anaemia.

The localisation of the intrinsic factor in the pylorus and duodenum of the pig was achieved by Meulengracht (1939) but Fox and Castle (1942) demonstrated that in man this factor was present mainly in the body of the human stomach so that the gastric lesion could be correlated with the absence of intrinsic factor. Coldhamer (1937) had shown that the gastric juice of patients with pernicious anaemia contained minute amounts of intrinsic factor but because of the achylia he suggested that the defect was mainly quantitative.

Intrinsic factor has since been identified as a mucopolypeptide (Williams *et al* 1954) present in the mucoprotein of gastric juice (Fatner *et al* 1954). It now seems likely that intrinsic factor combines with extrinsic factor (vitamin B<sub>12</sub> or cyanocobalamin isolated by Smith 1948 and Rickes *et al* 1948) and facilitates its absorption in the small intestine (Ross 1950). It is stored in the liver combined to a beta globulin (Pitney *et al* 1955) and then combined to an alpha globulin (Pitney *et al* 1954) it is transported in the plasma to the sites of utilisation. In conjunction with folic acid vitamin B<sub>12</sub> is thought to play a major role in the synthesis of nucleoprotein particularly at the sites of rapid cell proliferation such as the bone marrow. Vitamin B<sub>12</sub> deficiency occurring as a result of deficiency of intrinsic factor is followed by a decrease in the rate of cellular proliferation. The cells instead of dividing

and maturing become larger and remain immature, giving rise to megaloblasts

Vitamin  $B_{12}$  is stored in the liver in liberal amounts and its utilisation is slow. Thus when the absorption from the alimentary canal is retarded or ceases completely there is a delay in the development of overt pernicious anaemia or subacute combined degeneration of the cord. As stated previously this is demonstrated after total gastrectomy when it may take several years for megaloblastic anaemia to develop.

The relationship of vitamin  $B_{12}$  deficiency to the demyelinating peripheral neuritis and subacute combined degeneration of the cord is obscure. The lesions have been attributed to defective ribose nucleic acid synthesis affecting especially the long axones of nerve cells in the spinal cord (Nieweg *et al* 1954).

### Can pernicious anaemia be diagnosed in the presence of normal acid secretion and normal appearance of the gastric biopsy?

Mollin *et al* (1955) presented a case of pernicious anaemia without gastric atrophy. This case was initially diagnosed at the age of 13 months. The father of this patient subsequently developed an early pernicious anaemia which was associated with gastric atrophy. The mother had a brother who died of carcinoma of the stomach when aged 30 years. Moreover it is of considerable interest that the mother and father were cousins as consanguinity greatly enhances genetic influences. It was suggested that the early development of pernicious anaemia in the propositus was genetically determined, the patient being homozygous with respect to the faulty gene. It was also suggested that the early treatment of the case prevented the development of gastric atrophy and that gastric atrophy was in fact secondary to prolonged vitamin  $B_{12}$  deficiency which might be clinically silent for a long period. Similarly Harris Jones *et al* (1957) reported pernicious anaemia without gastric atrophy in a girl of 16 years.

Pernicious anaemia is very rare in children but Reisner *et al* (1951) found 5 cases in the literature in which hydrochloric acid was found in the gastric juice and they described 3 further cases in which there was hydrochloric acid in the gastric juice at some stage. This suggests that in the pernicious anaemia of

childhood achlorhydria is not so constant and that presumably the gastric atrophy is either absent or incomplete.

This finding is consistent with the concept of Mollin and his colleagues but it must be emphasized that such cases are at most rarely found—in classical pernicious anaemia the atrophic changes in the gastric mucosa are severe and achlorhydria is present.

Indeed we have not seen a case of pernicious anaemia without gross atrophy of the gastric mucosa and achlorhydria. Like Witts (1956) we wish to reserve judgment on the cases in this group. We do not consider they should be diagnosed as suffering from pernicious anaemia.

### Definition of "pernicious anaemia"

It is of interest that Davis and Brown (1953) defined pernicious anaemia as a megaloblastic anaemia with irreversible degeneration of the gastric mucosa. Witts (1956) stated the final argument for the priority of gastric atrophy in the development of pernicious anaemia is the complete failure to reverse the atrophy and achylia by vitamin B<sub>12</sub> or any other therapeutic agent. In our present state of knowledge we agree with Witts.

### Treatment of gastric atrophy, pernicious anaemia and subacute combined degeneration of the cord

Once the diagnosis of pernicious anaemia or subacute combined degeneration of the cord is established intramuscular injections of vitamin B<sub>12</sub> should be given. Before overt macrocytic anaemia or subacute combined degeneration of the cord appear there is a gradual depletion of the vitamin B<sub>12</sub> stores especially in the liver so that when treatment is begun these stores should be rapidly and fully replenished. It is wise to give 1000 micrograms of vitamin B<sub>12</sub> every second day for the first week and then weekly for 2 months. Iron also may be required to restore the normal blood picture. After 2 months liberal treatment with vitamin B<sub>12</sub> smaller doses of 100 micrograms can be given every 3–4 weeks to maintain adequately the erythrocyte count and haemoglobin level. A few patients appear to require 200 micrograms or more but once vitamin B<sub>12</sub> fails to control the anaemia other causes should be sought especially cancer of the stomach or bowel or chronic renal disease.

Should the initial anaemia be profound and the patient's life threatened and especially if there be an accompanying infection such as pneumonia then a blood transfusion can be life saving. The blood should be carefully matched and cross typed for transfusion reactions are frequently encountered in these patients. The use of packed cells to avoid overloading of the circulation with plasma should be considered. However most cases can be restored to health with liberal doses of vitamin B<sub>12</sub> and appropriate antibiotics without resorting to transfusion.

Throughout the present studies full doses of parental vitamin B<sub>12</sub> have been employed. Preparations for oral administration containing both B<sub>12</sub> and intrinsic factor in the form of animal (heterologous) mucosal extract have been disappointing in that their potency may wane after the passage of months or years. Schwartz and Meulengracht (1957) have attributed this fall in potency to the development of antibodies to heterologous extract. They demonstrated human (autologous) gastric mucosal extract to be still efficient when the animal extract was failing.

The treatment of patients with severe subacute combined degeneration of the cord with or without accompanying pernicious anaemia requires the same initial liberal dosage of vitamin B<sub>12</sub>. It has been claimed that with continued high dosage good results are obtained. All patients show improvement but in severe cases the recovery is incomplete. Moreover should the residual weakness be pronounced it may increase with advancing years probably due to failing blood supply from progressive atherosclerosis.

It has already been emphasised that the gastric lesion in classical pernicious anaemia and subacute combined degeneration of the cord is irreversible despite long continued therapy with liver preparations (Doig and Wood 1950, Wilkinson 1949) and with vitamin B<sub>12</sub> (Finckh and Wood 1953). Thus gastric biopsy is of value in reviewing the diagnosis in patients who have received some previous treatment with liver or vitamin B<sub>12</sub>. It is also of value in the diagnosis of pernicious anaemia as opposed to other types of megaloblastic anaemia in which severe gastritis or gastric atrophy may not occur namely those due to dietetic deficiency, intestinal malabsorption, liver disease and drugs such as I panu tin \* (Badenoch 1954).

Delay in the diagnosis and treatment of subacute combined degeneration of the cord may result either from failure to recognise

\* Epanutin (sodium phenytoin) Parke Davis

the significance of the peripheral dysaesthesiae which may precede other manifestations by many months or from a reluctance to diagnose this condition in the absence of the characteristic haematological abnormalities. Here gastric biopsy has been a most valuable aid in making an early diagnosis (Doig Motteram Robertson and Wood 19504). The serum vitamin B<sub>12</sub> estimations and radioactive vitamin B<sub>12</sub> absorption tests are also proving to be of considerable value.

Alcoholic peripheral neuritis associated with chronic atrophic gastritis and histamine fast achlorhydria may simulate early sub-acute combined degeneration of the cord.



## EPILOGUE

### And what of the future?

In the light of experience in our unit and elsewhere will gastritis become a widely accepted malady sympathetically managed by the attending physician? Will the atomic age with increasing exposure to radiation produce more sufferers from chronic atrophic gastritis? Will this in turn lead to more cases of pernicious anaemia? And more cases of gastric cancer? And fewer cases of duodenal ulcer? What will be the future of alcoholism—will it be successfully controlled by social reform? And what will be the final assessment of the effect on the gastric mucosa of the stress reactions of Selye and of Wolf and Wolff?

Let us look into the future and encourage our successors to study man and his way of life in this changing world. It will be of particular interest to gastroenterologists to observe the impact of man's way of life on the structure and function of the gastric mucosa.

## REFERENCES

- ADDISON T (1849) Anaemia—disease of the supra renal capsules  
*Lond med Gaz* 8 517
- ALTHUSEN T L, DOIG R K, WEIDEN S, MOTTERAM R, TURNER C V and MOORE A (1951) Haemochromatosis *A M A Arch intern Med* 88 553
- BADENOCH J (1954) Section of experimental medicine and therapeutics discussion gastric biopsy and the investigation of the megaloblastic anaemias. The use of labelled vitamin B<sub>12</sub> and gastric biopsy in the investigation of anaemia *Proc roy Soc Med* 47 426
- BADENOCH J and RICHARDS W C D (1953) Gastric lesion in anaemia with particular reference to biopsy *Gastroenterologia* 79 329
- BADENOCH J and CALLENDER S T E (1954) Iron metabolism in steatorrhea use of radioactive iron in studies of absorption and utilization *Blood* 9 123
- BASSLER (1909) Some medical considerations in the treatment of gastric ulcer with special reference to the use of belladonna and X ray therapy *Med Times* 37 259
- BEAUMONT W (1833) *Experiments and observations on the gastric juice and the physiology of digestion* F P Allen Plattsburgh New York
- BENEDICT E H (1948) An operating gastroscope *Gastroenterology* 2 281
- BENSLEY H H (1928) *The gastric glands in Special Cytology* 1 137 E V Cowdry New York
- BOOTH C C and MOLLER D L (1956) Plasma tissue and urinary radioactivity after oral administration of <sup>54</sup>Co labelled vitamin B<sub>12</sub> *Brit J Haematology* 2 223
- BOWIE D J and VINEBERG A M (1935) Selective action of histamine and effect of prolonged vagal stimulation on cells of gastric glands in the dog *Quart J exp Physiol* 25 247
- BROUSSAIS F J V (1831) *History of chronic phlegmasiae or inflammations* Translated from the French of the 4th ed by Isaac Hays and R Eglesfeld Griffith Carey and Lear Philadelphia
- BROWN G and WOOD I J (1955) Antroduodenectomy and X ray irradiation in the treatment of duodenal ulcer a progress report *Aust N Z J Surg* 25 260
- (1956) Treatment of chronic duodenal ulcer by antroduodenectomy and X irradiation *Lancet* 2 169

- BROWN G SCOTT R K HOLMAN W P WOOD I J FINCKH E S WEIDEN S and DAVIS P (1952) Antroduodenectomy and X ray irradiation in the treatment of duodenal ulcer *Lancet* 2, 1145
- CASTLE W B TOWNSEND W C HEATH C W and STRAUSS M H (1929) Observations on etiologic relationship of achylia gastrica to pernicious anaemia I *Amer J med Sci* 178 748
- COGHILL N F and WILLIAMS A W (1955) Gastric biopsy with a modified Australian instrument *Brit med J* 2 1111
- COHN A and VON MERING J (1886) Die Sauren des gesunden und kranken Magens *Dtsch Arch Klin Med* 39 233
- COX A J (1943) The stomach in pernicious anaemia *Amer J Path* 19 491
- (1952) Stomach size and its relation to chronic peptic ulcer *Arch Path* 54 407
- CROZIER R E MIDDLETON M and ROSS J R (1956) Clinical application of gastric cytology *New Engl J Med* 255 1128
- D ALMEIDA C (1948) Foreign letter *J Amer med Ass* 136 84
- DAVIDSON W M B and MARKSON J L (1955) The gastric mucosa in iron deficiency anaemia *Lancet* 2 639
- DAVIDSON S (1957) Clinical picture of pernicious anaemia prior to introduction of liver therapy in 19-6 and in Edinburgh subsequent to 1944 *Brit med J* 2 241
- DAVIS L J and BROWN A (1953) *The megaloblastic anaemias* Blackwell Oxford
- DOIG R K (1949) Some observations on chronic gastritis with reference to findings of gastric biopsy *Proc roy Aust Coll Physicians* 4 90
- (1954) Section of experimental medicine and therapeutics Discussion gastric biopsy and the investigation of the megaloblastic anaemias Gastric biopsy and gastritis *Proc roy Soc Med* 47 243
- DOIG R K FUNDEN J F and WEIDEN S (1951) Serial gastric biopsy studies in a case of duodenal ulcer treated by deep X ray therapy *Med J Aust* 1 828
- DOIG R K and MOTTERAM R (1950) The histology of gastritis *Aust N Z J Surg* 19 96
- DOIG R K and WOOD I J (1950) Gastric biopsy in pernicious anaemia *Med J Aust* 2 565
- DOIG R K MOTTERAM R ROBERTSON F and WOOD I J (1950A) Early diagnosis of subacute combined degeneration of the cord Value of gastric biopsy *Lancet* 2 836
- DOIG R K MOTTERAM R WEIDEN S and WOOD I J (1950B) Achlorhydria in young adults A study of the gastric mucosa obtained by biopsy from a volunteer *Lancet* 1 948

- DOIG R K and WOOD I J (1952A) Gastritis a study of 112 cases diagnosed by gastric biopsy *Med J Aust* 1 593
- (1952B) *Modern trends in gastroenterology* Butterworth and Co London 352
- (1952C) Gastric and duodenal ulcer a survey of 141 cases *Med J Aust* 1 60.
- DLBACH R CALLENDER S T F and MOORE C V (1948) Studies in iron transportation and metabolism VI Absorption of radio active iron in patients with fever and with anaemias of varied etiology *Blood* 3 5-6
- EDWARDS F C and EDWARDS J H (1956) Tea drinking and gastritis *Lancet* 2 543
- EINHORN M (1894) Clinical observations on erosions of the stomach and their treatment *Med Rec* 45 780
- EPSTEIN I S MACKAY M A and TURNER C V (1950) Control of malnutrition *Med J Aust* 1 799
- FABER K (1935) *Gastritis and its consequences* Oxford Medical Publications
- FABER K and BLOCH C E (1900) Über die Pathologische Veränderungen am Digestions tractus bei der Perniciösen Anämie und über die sogenannte Dermatrophy *Zeitschrift für Klinische Med* V 40 98
- FAIRLEY H F TURNER C V MACKAY M A and JOHNS H A (1955) Atrophic gastritis a five year survey of 32 cases proven by gastric biopsy *Med J Aust* 2 1085
- FENWICK S (1870) On atrophy of the stomach *Lancet* 2 78
- (1880) *On atrophy of the stomach and the nervous affections of the digestive organs* London 1880
- FIEBER S S (1955) Hypertrophic gastritis Report of two cases and analysis of fifty pathologically verified cases from the literature *Gastroenterology* 28 39
- FINCKH E S DALE B G JOSKE R A and SAINT E G (1955) Studies in chronic alcoholism 3 Laboratory studies *Med J Aust* 1 738
- FINCKH E S and WOOD I J (1953) Failure of vitamin B<sub>12</sub> therapy to improve gastric atrophy in pernicious anaemia report of a case studied by serial gastric biopsies *Gastroenterology* 25 48
- FLINT A (1860) *Amer med Times* 1 181
- FOX H J and CASTLE W B (1942) Observations on the etiologic relationship of achylia gastrica to pernicious anaemia IX Difference in site of secretion of intrinsic factor in the hog and in the human stomach *Amer J med Sci* 203 18
- FLANDER J F and WEIDEN S (1952) The correlation between test meal findings and the histology of the stomach as shown by gastric biopsy *Med J Aust* 1 600

- GOLDHAMER S M (1937) Gastric juice in patients with pernicious anaemia in induced remission *Amer J med Sci* 193 23
- GRAHAM R M and RHEAULT M H (1954) Characteristic cellular changes in epithelial cells in pernicious anaemia *J Lab clin Med* 43 235
- GRASBECK R (1956) Studies of the vitamin B<sub>12</sub> binding principle and other biocolloids of human gastric juice *Acta med scand Suppl* 314 1
- GRAY S RAVISAY C REIFENSTEIN R W and BENSON J A (1953) The significance of hormonal factors in the pathogenesis of peptic ulcer *Gastroenterology* 25 156
- GUNTER G S (1950) A histological investigation of the healing of acute gastric ulceration in the cat *Gastroenterology* 15, 708
- HAGEDORN A B (1957) Intramuscularly administered iron in the treatment of iron deficiency anaemia *Amer med Ass J* 64 1642
- HARRIS JONES J N SWAN H T and TUDHOPE G R (1957) Pernicious anaemia without gastric atrophy and in the presence of free hydrochloric acid. Report of a case *Blood* 12 461
- HARVEY J C (1956) The vitamin B<sub>12</sub> deficiency state engendered by total gastrectomy *Surgery* 40, 977
- HAWKSLEY J (1939) A note on the occurrence of pieces of sloughed gastric mucous membrane in aspirated gastric juice *J Path Bact* 49 585
- HAWKSLEY J C and COORAY G H (1948) Observations on fragments of gastric mucous membrane found in aspirated resting gastric juice *J Path Bact* 60 333
- HEBBEL R (1949) The topography of chronic gastritis in otherwise normal stomachs *Amer J Path* 25 125
- HOAR C S and BROWNING J R (1956) Plasma pepsinogen in peptic ulcer and other gastric disorders a clinical and laboratory evaluation *New Engl J Med* 255 4
- HOLMAN W P and LEWIS R A (1941) Some observations of the effects of selective irradiation of the stomach in cases of chronic non obstructive duodenal ulcer with hyperchlorhydria *Med J Aust* 2 734
- HOLMES J M (1956) Cerebral manifestations of B<sub>12</sub> deficiency *Brit med J* 2 1394
- HURLEY T and BAZELEY V (1957) Absorption of penicillin from the human stomach *Unpublished*
- JACKSON C (1907) Gastroscopy *Med Rec* 71, 549
- JACKSON C and JACKSON C L (1935) Percoral gastroscopy including examination of supradiaphragmatic stomach *J Amer med Ass* 104 269

- JOSKE R A FINCKH E S and WOOD I J (1955) Gastric biopsy: a study of 1000 consecutive successful gastric biopsies. *Quart J Med (New Series)* 24: 69
- JOSKE R A and TURNER C N (1952) Studies in chronic alcoholism. 1. The clinical findings in 78 cases of chronic alcoholism. *Med J Aust* 1: 3
- JOSKE R A McALLISTER J C and PRANKERD T A J (1956) Isotope investigations of red cell production and destruction in chronic renal disease. *Clin Sci* 25: 511
- KATZ D and SELENICK S (1956) Massive pneumoperitoneum and pneumoretroperitoneum after gastroscopy. Report of a case and review of the literature. *Amer J dig Dis* 1, 12: 512
- KAY A W (1953) Effect of large doses of histamine on gastric secretion of HCl. An augmented histamine test. *Brit med J* 2, 77
- KENAMORE B (1940) Biopsy forceps for flexible gastroscope. *Amer J dig Dis* 7: 539
- KENAMORE B SCHEFF H and WOMACK A (1946) Study of gastric lesions by means of biopsy specimens removed endoscopically. *A M A Arch Surg (Chicago)* 52: 50
- KOJETZKY G F (1928) Die Entzündung des Magens. In Henle Lubarsch *Handbuch d Pathol Anatomie* 4 Heft 2 (Berlin Springer)
- LANDBOE CHRISTENSEN E BERG L and CASTLE W B (1952) Observations of the etiologic relationship of achylia gastrica to pernicious anaemia. IV. Intrinsic factor activity of desiccated and defatted human fundus duodenum and jejunum. *Amer J med Sci* 224: 1
- LATNER A L MERRILLS R J and RAINE I C D P (1954) Isolation of Castle's intrinsic factor. *Lancet* 1: 497
- (1954) Further observations on the isolation and chemical composition of Castle's intrinsic factor. *Biochem J* 57: Proc Sect 19
- LEVY E CLAYMAN C B PALMER W C and KIRKNER J B (1957) Observations on the value of gastric irradiation in the treatment of duodenal ulcer. *Gastroenterology* 32: 42
- LEVY A H and LEVINE S (1956) Pepsin excretion studies in clinical surgery. *Gastroenterology* 30: 270
- LIVERSIDGE L A BERYNE G M and EMERY E W (1957) Radioactive B<sub>12</sub> in the diagnosis of neurological disorders. *Lancet* 1: 294
- MCLINTYRE P A SACHS M V KREVANS J R and CONLEY C L (1956) Pathogenesis and treatment of macrocytic anaemia. Information obtained with radioactive vitamin B<sub>12</sub>. *A M A Arch intern Med* 96: 541

- MAGNUS H A (1937) Observations on the presence of intestinal epithelium in the gastric mucosa *J Path Bact* 44 389
- (1952) Gastritis *Modern trends in gastroenterology* Butterworth London
- MAGNUS H A and UNGLEY C C (1938) The gastric lesion in pernicious anaemia *Lancet* i 420
- MARASON J L and DAVIDSON W M B (1956) Gastric biopsy in the megaloblastic anaemias *Scot med surg J* 1, 259
- MASSEY B W and RUBIN C E (1954) The stomach in pernicious anaemia A cytological study *Amer J med Sci* 227, 481
- MATZNER M J RAAB A P and SPEAR P W (1951) Benign giant gastric rugae complicated by submucosal gastric carcinoma *Gastroenterology* 18 296
- MAXIMOW A A and BLOOM W (1948) A textbook of histology W B Saunders Co Philadelphia and London
- MEULENGRACHT E (1939) Histologic investigation of pyloric gland organ in pernicious anaemia *Amer J med Sci* 197, 201
- MOLLIN D L BAKER S G and DONIACH G (1955) Addisonian pernicious anaemia without gastric atrophy in a young man *Brit J Haematology* 1 278
- MOORE C V (1955) The importance of nutritional factors in the pathogenesis of iron deficiency anaemia *Amer J clin Nutrition* 3 3
- MOSBECH J (1953) Heredity in pernicious anaemia Copenhagen
- MOTTERAM R (1951) A biopsy study of chronic gastritis and gastric atrophy *J Path Bact* 63 389
- NIEWEG H O FABER J G DE VRIES J A and KROESE W F S (1954) The relationship of vitamin B<sub>12</sub> and folic acid in megaloblastic anaemias *J Lab clin Med* 44 118
- NYFELDT A and VIMTRUP B (1932) Gastritis acuta bei diphtheria faucium *Acta med scand* 78 447
- PALMER E D (1950) Gastric mucosal biopsy findings correlated with gastroscopic diagnoses Preliminary report based on fifty patients *Amer J med Sci* 219 648
- (1951) The morphologic consequences of acute exogenous (staphylococcic) gastroenteritis on the gastric mucosa *Gastroenterology* 19 46
- (1954) Gastritis—A revaluation *Medicine* 33 199
- (1954) State of the gastric mucosa of elderly persons without upper gastrointestinal symptoms *J Amer Geriatrics Soc* 2 171
- PALMER E D and SMITH V M (1953) Submucosal gastric hematomia Complication of vacuum tube biopsy technic *Review of Gastroenterology* 20 330

- PALUMBO L T RUGTIV G M and CROCK H R (1951) Giant hypertrophic gastritis Its surgical and pathologic significance *Ann Surg* 134 259
- PARSONS P J TURNER C N and WOOD I J (1948) Some causes of malnutrition *Proc roy Aust Coll Physicians* 3 107
- PERRY J F YONEHIRO E G YAP M ROOT H D and WANGENSTEEN O H (1956) Digestive action of human gastric juice *Proc Soc exp Biol NY* 92 237
- PITNEY W R BEARD M F and VAN LOON E J (1954) Observations on the bound form of vitamin B<sub>12</sub> in human serum *J biol Chem* 207 143
- (1955) The vitamin B<sub>12</sub> content of electrophoretic fractions of liver homogenates *J biol Chem* 212 117
- PLENCK H (1932) Der Magen In Moellendorf *Handbuch d mikr Anatomie d Menschen* 5 Teil 2 (Berlin Springer)
- RECHTSCHAFFEN J S and WEINGARTEN M (1957) Gastric acidity determination procedures—Comparison of a tubeless method with azure A resin compound and an intubation method with histamine *J Amer med Ass* 164 1467
- REISNER E H WOLFF J A MCKAY R J and DOYLE M F (1951) Juvenile pernicious anaemia *Pediatrics* 8 88
- RICKES E L BRINK N G KONINSZY F R WOOD T R and FOLDERS H (1948) Crystalline vitamin B<sub>12</sub> *Science* 107, 396
- RICKETTS W E KIRSNER J B and PALMER W L (1947) Large otherwise normal gastric rugae simulating tumor of the stomach A report of three cases *Gastroenterology* 8 123
- RICKETTS W E PALMER W L KIRSNER J B and HAMANN A (1948) Radiation therapy in peptic ulcer An analysis of results *Gastroenterology* 11 789
- ROBERTSON E G WOOD I J and JOCKE R A (1955) Gastric atrophy with subsequent pernicious anaemia and pronounced subacute combined degeneration of the cord *Lancet* 2 69
- ROSS G I M (1950) Vitamin B<sub>12</sub> assay in body fluids *Nature* 166 70
- RUBIN C E GOLDGRABER M H and SMITH C (1953) An improved gastric biopsy tube—Its manufacture use and investigative value *Gastroenterology* 25 31
- SANDERS G H and MEGRAY P M (1941) Pseudogastritis of operative origin *Ann Surg* 114 986
- SCHINDLER R (1952) Die Diagnostische Bedeutung der Gastroskopie *Munchen med Wchnschr* 69 535
- (1937) *Gastroscoy The endoscopic study of gastric pathology* University of Chicago Press Chicago Ill
- (1947) *Gastritis* W Heinemann London



- MAGNUS H A (1937) Observations on the presence of intestinal epithelium in the gastric mucosa *J Path Bact* 44 389
- (1952) Gastritis *Modern trends in gastroenterology* Butterworth London
- MAGNUS H A and UNGLEY C C (1938) The gastric lesion in pernicious anaemia *Lancet* i 420
- MARKSON J L and DAVIDSON W M B (1956) Gastric biopsy in the megaloblastic anaemias *Scot med surg J* i 259
- MASSEY H W and RUBIN C E (1954) The stomach in pernicious anaemia A cytological study *Amer J med Sci* 227 481
- MATZNER M J RAAB A P and SPEAR P W (1951) Benign giant gastric rugae complicated by submucosal gastric carcinoma *Gastroenterology* 18 296
- MAXIMOW A A and BLOOM W (1948) A textbook of histology W B Saunders Co Philadelphia and London
- MEULENGRACHT E (1939) Histologic investigation of pyloric gland organ in pernicious anaemia *Amer J med Sci* 197 201
- MOLLIN D L BAKER S G and DONNICH G (1955) Addisonian pernicious anaemia without gastric atrophy in a young man *Brit J Haematology* 1, 278
- MOORE C V (1955) The importance of nutritional factors in the pathogenesis of iron deficiency anaemia *Amer J clin Nutrition* 3 3
- MOSBECH J (1953) Heredity in pernicious anaemia Copenhagen
- MOTTERAM R (1951) A biopsy study of chronic gastritis and gastric atrophy *J Path Bact* 63 389
- NIEWEG H O FABER J G DE VRIES J A and KROESE W F S (1954) The relationship of vitamin B<sub>12</sub> and folic acid in megaloblastic anaemias *J Lab clin Med* 44 118
- NYFELDT A and VIMTRUP B (1932) Gastritis acuta bei diphtheria faucium *Acta med scand* 78 447
- PALMER E D (1950) Gastric mucosal biopsy findings correlated with gastroscopic diagnoses Preliminary report based on fifty patients *Amer J med Sci* 219 648
- (1951) The morphologic consequences of acute exogenous (staphylococcal) gastroenteritis on the gastric mucosa *Gastroenterology* 19 462
- (1954) Gastritis—A revaluation *Medicine* 33 199
- (1954) State of the gastric mucosa of elderly persons without upper gastrointestinal symptoms *J Amer Geriatrics Soc* 2 171
- PALMER E D and SMITH V M (1953) Submucosal gastric hematoma Complication of vacuum tube biopsy technic *Review of Gastroenterology* 20 330

- WILLIAMS W L ELLENBOGEN L and ESPOSITO R G (1954) Preparation of highly purified intrinsic factor (*Proc Soc exper Biol* 87 400)
- WITTS L J (1956) Anaemia and the alimentary tract The relationship between changes in the alimentary tract and deficiencies of iron folic acid and vitamin B<sub>12</sub> (*Royal Coll Phys Ed Publications* No 7 37)
- WOLF E and WOLFF H G (1943) *Human gastric function An experimental study of a man and his stomach* Oxford University Press London
- WOLFF H G WOLF S GARE W J HOLMES T F STEVENSON I STRAUB L GOODELL H and SETON P (1948) Changes in form and function of mucous membranes occurring as part of preventive reaction patterns in man during periods of life stress and emotional conflict (*Transactions 4th Amer Phys* 61, 313)
- WOOD I J (1951) Value of gastric biopsy in the study of chronic gastritis and pernicious anaemia (*Brit med j* 2, 8-3)
- WOOD I J DOIG R K MOTTERAM R and HUGHES A (1949A) Gastric biopsy Report on fifty five biopsies using a new flexible gastric biopsy tube (*Lancet* 1 19)
- WOOD I J DOIG R K MOTTERAM R WEIDEN E and MOORE A (1949B) The relationship between the secretions of the gastric mucosa and its morphology as shown by biopsy specimens (*Gastroenterology* 12 949)

- SCHINDLER R NECHIEFS H and GOLD H (1939) Surgical gastritis. A study on the genesis of gastritis found in resected stomachs with particular reference to the so called Antral Gastritis associated with ulcer *Surg Gynec Obstet* 69 281
- SCHINDLER R and ORTMAYER M (1942) Histopathology of chronic gastritis *Am J dig Dis* 9 411
- SCHWARTZ M LOUS P and MEULENGRACHT L (1957) Reduced effect of heterologous intrinsic factor after prolonged oral treatment in pernicious anaemia *Lancet* 1, 751
- SHINER M and DONACH I (1957) A study of X ray negative dyspepsia with reference to histologic changes in the gastric mucosa *Gastroenterology* 32 313
- SMITH E L (1948) Purification of anti pernicious anaemia factors from liver *Nature* 161 638
- STEIGMANN F HYMAN S and KANNAPEL W L (1957) Large gastric rugae Benign or malignant *Gastroenterology* 32, 7-
- SWALM W A JACKSON C L and MORRISON L (1936) Correlation of clinical and gastroscopic findings in chronic gastritis *Rev Gastroenterol* 3 219
- TFYTER E C (Jnr) LEGERTON C W REEVE R J SMITH A G and RUFFIN J M (1953) Coexistent carcinoma of the stomach and hypertrophic gastritis *Gastroenterology* 24 579
- THOMSEN F (1925) Études sur l'achylie neurogene et cellulaire I *Acta med scand* 61, 377
- TOMENIUS J (1952) A new instrument for gastric biopsies under visual control *Gastroenterology* 21 544
- VICTOR M and LEAR A (1956) Subacute combined degeneration of the spinal cord Current concepts of the disease process Value of serum B<sub>12</sub> determinations in clarifying some of the problems *Amer J Med* 20 896
- VIDEBAEK A and MOSBECH J (1954) The aetiology of gastric carcinoma elucidated by a study of 30- pedigrees *Acta med Scand* 149 137
- VIMTRUP B (1929) Two lectures on gastritis *Bibliothek f Laeger* 212 119
- WEIDEN S (1949) Some observations on the secretion of the stomach with special reference to mucus *Med J Aust* 2 81
- WELBOURN R B NELSON M G and ZACHARIAS F J (1956) Megaloblastic anaemia following gastric resection *Brit J Sur* 43 422
- WILKINSON J F (1949) Megalocytic anaemias *Lancet* 1 249
- WILLIAMS A W (1950) The stomach of the recently deceased *Brit med J* 1 102
- (1956) Effects of alcohol on gastric mucosa *Brit med J* 1 256

- WILLIAMS W L ELLENBOGEN L and E PO-ITO R G (1954) Preparation of highly purified intrinsic factor *Proc Soc exper Biol* 87, 400
- WITTS L J (1956) Anaemia and the alimentary tract The relationship between changes in the alimentary tract and deficiencies of iron folic acid and vitamin B<sub>12</sub> *Royal Coll Phys Ed Publications* No 7 37
- WOLF S and WOLFF H G (1943) *Human gastric function An experimental study of a man and his stomach* Oxford University Press London
- WOLFF H G WOLF S GARE W J HOLMES T E STEVENSON I STRAUB L GOODELL H and SFTON P (1948) Changes in form and function of mucous membranes occurring as part of preventive reaction patterns in man during periods of life stress and emotional conflict *Transactions Am Amer Phys* 61 313
- WOOD I J (1951) Value of gastric biopsy in the study of chronic gastritis and pernicious anaemia *Brit med J* 2 823
- WOOD I J DOIG R H NOTTERAM R and HUGHES A (1949A) Gastric biopsy Report on fifty five biopsies using a new flexible gastric biopsy tube *Lancet* 1, 18
- WOOD I J DOIG R H NOTTERAM R WEIDEN E and MOORE A (1949B) The relationship between the secretions of the gastric mucosa and its morphology as shown by biopsy specimens *Gastroenterology* 12 949

- SCHINDLER R, NECHELES H and GOLD R (1939) Surgical gastritis. A study on the genesis of gastritis found in resected stomachs with particular reference to the so called Antral Gastritis associated with ulcer. *Surg Gynec Obstet* 69 281
- SCHINDLER R and ORTMAYER M (1942) Histopathology of chronic gastritis. *Am J dig Dis* 9 411
- SCHWARTZ M, LOIS P and MELLENGRACHT F (1957) Reduced effect of heterologous intrinsic factor after prolonged oral treatment in pernicious anaemia. *Lancet* 1, 751
- SHINER M and DONACH I (1957) A study of X ray negative dyspepsia with reference to histologic changes in the gastric mucosa. *Gastroenterology* 32 313
- SMITH E L (1948) Purification of anti pernicious anaemia factors from liver. *Nature* 162, 638
- STEIGMANN F, HYMAN S and KANNAPEL W L (1957) Large gastric rugae. Benign or malignant. *Gastroenterology* 32 7-
- SWALM W A, JACKSON C L and MORRISON I (1936) Correlation of clinical and gastroscopic findings in chronic gastritis. *Rev Gastroenterol* 3 219
- TEYTER E C (Jnr), LEGERTON C W, REEVES R J, SMITH A G and RUFFIN J M (1953) Coexistent carcinoma of the stomach and hypertrophic gastritis. *Gastroenterology* 24, 579
- THOMSEN E (1925) Études sur l'achylie neurogene et cellulaire. I. *Acta med scand* 61 377
- TOMENIUS J (1952) A new instrument for gastric biopsies under visual control. *Gastroenterology* 21 544
- VICTOR M and LEAR A (1956) Subacute combined degeneration of the spinal cord. Current concepts of the disease process. Value of serum B<sub>12</sub> determinations in clarifying some of the problems. *Amer J Med* 20 896
- VIDEBÆK A and MOSBECH J (1954) The aetiology of gastric carcinoma elucidated by a study of 302 pedigrees. *Acta med Scand* 149 137
- VINTRUP B (1929) Two lectures on gastritis. *Bibliothek f Laeger* 212 119
- WEIDEN S (1949) Some observations on the secretion of the stomach with special reference to mucus. *Med J Aust* 2 81
- WELBOURN R B, NELSON M G and ZACHARIAS F J (1956) Megaloblastic anaemia following gastric resection. *Brit J Sur* 43 4-2
- WILKINSON J F (1949) Megalocytic anaemias. *Lancet* 1 49
- WILLIAMS A W (1950) The stomach of the recently deceased. *Brit med J* 1 102
- (1956) Effects of alcohol on gastric mucosa. *Brit med J* 1 56

- Gastric atrophy—contd**  
 microscopic appearance 57 58  
 preceding pernicious anaemia 61  
 relation to pernicious anaemia  
   57 64 65 66 67 68  
 without pernicious anaemia 41  
 as a primary process 57 65  
 symptomatology 61  
 treatment of 69  
 lack of response to vitamin B<sub>12</sub>  
   15 60 70
- Gastric biopsy**  
 in achlorhydria 15  
 in alcoholism 22 39 41  
 artefacts 14  
 complications 16  
 contraindications 15  
 efficiency of flexible tube 10  
 structure of flexible tube 7 9  
 in gastric atrophy 57 58 59 60  
 relation to gastric secretion 66  
   67  
 in acute gastritis 19 20 21 22  
   39  
 in chronic gastritis 24  
 serial in chronic gastritis 47  
 in diffuse giant hypertrophic  
   gastritis 54  
 in radiation gastritis 42 44  
 relation to gastroscopy 5  
 history 6  
 indications 15  
 in megaloblastic anaemia 15 63  
   70  
 normal appearance 13 14  
 correlation with pepsin secretion  
   15  
 in pernicious anaemia 59 63 65  
   68  
 in diagnosis of pernicious anaemia  
   and subacute combined de-  
   generation of the cord 71  
 preparation of specimen 11 12  
   13  
 sampling error 11  
 staining of specimen 11  
 technique using flexible tube 7  
   9 10  
 technique using Benedict operat-  
   ing gastroscop 7  
 correlation with test meal 3 4  
 its value 1  
 in Witten's hypochloric anaemia  
   49
- Gastric fistula** 18
- Gastric mucosa**  
 effect of age on 38  
 effect of alcohol 2  
 argentaffine cells in 14  
 artefacts 14  
 changes after death and surgery  
   1  
 early fixation 1 18  
 goblet cells in 13  
 mitoses in 14  
 normal appearance in biopsy 13  
 secreting bulk 45
- Gastric secretion** *see also* Achlor-  
 hydria  
 effect of age 38  
 relation to gastric biopsy 66 67  
 in chronic gastritis 46 47 48  
 mucus content in chronic gastritis  
   and gastric atrophy 67  
 normal in possible case of per-  
   nicious anaemia 68 69  
 in peptic ulcer 45 46  
 following X-ray irradiation 42  
   44
- Gastritis acute**  
 — in alcoholism 22 39  
 — causes 18 21 22 23  
 — gastric biopsy in 19 20 21  
   22 39  
 — gastroscopic appearances 21  
   22  
 — gastroscopy in 5  
 — history 18  
 — macroscopic appearance 18  
 — microscopic appearance 19  
   20  
 — perforation in 20  
 — regeneration in 20  
 — due to staphylococcal entero-  
   toxin 21 23  
 — staphylococcal gastroenteritis  
   21 23 39  
 — symptomatology 20 21  
 — treatment 23
- Gastritis chronic**  
 — achlorhydria in 36  
 — acute exacerbations in 27  
 — aetiology 37  
 — age of onset 36  
 — incidence with age 37 38  
 — alcoholism and 22 37 39 41  
 — relationship to cancer 36 46  
   47 48  
 — association with chronic de-  
   bilitating disease 36 37 39

# INDEX

- Achlorhydria**  
 age and 38  
 associated with alcoholism 71  
 in cancer 46  
 familial incidence of 66  
 in gastric atrophy 59 66  
 value of gastric biopsy in 15  
 in chronic gastritis 36  
 in healthy subjects 66  
 iron deficiency in 48  
 and peripheral neuritis 71  
 in pernicious anaemia and sub  
 acute combined degeneration  
 of the cord 59 63 67 69
- Acute gastritis** *see* Gastritis acute
- Alcohol**  
 effect on diet ~ 37 39 41  
 effect on gastric mucosa ~ 22  
 39 41  
 and acute gastritis 2~ 39  
 and chronic gastritis 2~ 37 39  
 41  
 and hepatomegaly 2  
 peripheral neuritis caused by 71
- Anaemia** *see also* Pernicious anaemia  
 carcinoma of colon causing 51  
 chronic nephritis causing 51  
 macrocytic 51  
 Wirtz hypochromic 48  
 — — gastritis in 48 49 50 51  
 — — gastric biopsy in 49  
 — — as precursor of cancer 48  
 — — vitamin B<sub>12</sub> absorption in  
 48 49
- Atrophic gastritis** *see* Gastritis  
 chronic
- Atrophy** *see* Gastric atrophy
- Biopsy** *see* Gastric biopsy
- Cancer of the stomach** 2 3  
 achlorhydria in 46  
 dyspepsia in 47  
 familial incidence 65 66  
 relationship to gastritis 36 46  
 47 48
- Cancer of the stomach—contd**  
 resembling chronic gastritis 5  
 zonal gastritis in relation to 47  
 relationship to pernicious  
 anaemia 66  
 X rays as possible cause of 48
- Carcinoma** of colon causing  
 anaemia 51
- Clinical history** 2
- Cytology of gastric washings** 17  
 in pernicious anaemia 60
- Diarrhoea in chronic gastritis** 36
- Diet**  
 in alcoholism 37 39 41  
 in cancer 47  
 in chronic gastritis 33 37 38  
 in clinical history ~
- Diphtheria** acute gastritis in ~3
- Dyspepsia**  
 age of onset in gastritis 36  
 in cancer 47  
 caused by gastritis 1 2  
 in chronic gastritis 33 36 51  
 in radiation gastritis 44  
 in gastric atrophy 61 6~  
 in pernicious anaemia 61 6  
 treatment with hydrochloric acid  
 33
- Flexible gastric biopsy tube** *see*  
 Gastric biopsy
- Gastrectomy** cause of vitamin B<sub>12</sub>  
 deficiency 6~ 68
- Gastric atrophy** 57 *see also* Per  
 nicious anaemia  
 achlorhydria in 59  
 aetiology of 64 65 66  
 dyspepsia in 61 6~  
 relation to chronic gastritis ~5  
 33 59 61 65 66  
 genetic influence in 65 66 68  
 history 57  
 macroscopic appearance 57

**Gastric atrophy—*contd***

- microscopic appearance 57 58
- preceding pernicious anaemia 61
- relation to pernicious anaemia 57 64 65 66 67 68
- without pernicious anaemia 41
- as a primary process 57 65
- symptomatology 61
- treatment of 69
- lack of response to vitamin B<sub>12</sub> 15 60 70

**Gastric biopsy**

- in achlorhydria 15
- in alcoholism 2- 37 41
- artefacts 14
- complications 16
- contraindications 15
- efficiency of flexible tube 10
- structure of flexible tube 7 9
- in gastric atrophy 57 58 59 60
- relation to gastric secretion 66 67
- in acute gastritis 19 0 21 -2 39
- in chronic gastritis 24
- serial in chronic gastritis 47
- in diffuse giant hypertrophic gastritis 54
- in radiation gastritis 4- 44
- relation to gastroscopy 5
- history 6 7
- indications 15
- in megaloblastic anaemia 15 63 70 71
- normal appearance 13 14
- correlation with pepsin secretion 15
- in pernicious anaemia 59 63 65 68
- in diagnosis of pernicious anaemia and subacute combined degeneration of the cord 71
- preparation of specimen 11 1 13
- sampling error 11
- staining of specimen 11
- technique using flexible tube 7 9 10
- technique using Benedict operation, gastroscope 7
- correlation with test meal 3 4
- its value 1
- in Witt hypochromic anaemia 49

**Gastric fistula 18****Gastric mucosa**

- effect of age on 38
- effect of alcohol 2
- argentaffine cells in 14
- artefacts 14
- changes after death and surgery 1
- early fixation 1 18
- goblet cells in 13
- mitoses in 14
- normal appearance in biopsy 13
- secreting bulk 45

**Gastric secretion *see also* Achlorhydria**

- effect of age 38
- relation to gastric biopsy 66 67
- in chronic gastritis 46 47 48
- mucus content in chronic gastritis and gastric atrophy 67
- normal in possible case of pernicious anaemia 68 69
- in peptic ulcer 45 46
- following X ray irradiation 4- 44

**Gastritis acute**

- in alcoholism 2- 39
- causes 18 21 22 23
- gastric biopsy in 19 -0 21 -2 39
- gastroscopic appearances -1 -2
- gastroscopy in 5
- history 18
- macroscopic appearance 16
- microscopic appearance 19 -0
- perforation in 20
- regeneration in 20
- due to staphylococcal enterotoxin -1 -3
- staphylococcal gastroenteritis 21 23 39
- symptomatology -0 -1
- treatment -3

**Gastritis chronic**

- achlorhydria in 36
- acute exacerbations in -7
- aetiology 37
- age of onset 36
- incidence with age 37 38
- alcoholism and -2 37 37 41
- relationship to cancer 36 46 47 48
- association with chronic debilitating disease 37 38 39



- Gastritis chronic classification 45
- diarrhoea in 36
  - diet in 33 37 38
  - distribution of 25
  - dyspepsia in 33 36 51
  - gastric atrophy and 25 33 59 61 65 66
  - serial gastric biopsy in 47
  - gastric biopsy in 24 47
  - atrophic gastritis 25 29 33 36 41 42 45 47 48
  - relation to diffuse giant hypertrophic gastritis 24 25 53
  - superficial gastritis 25 27 36 38 39 41 45
  - gastroscopy in 5
  - association with haemochromatosis 39
  - haemorrhage in 33 36 50
  - history 24
  - intestinal metaplasia in 29
  - resemblance to malignancy 52
  - macroscopic appearance 25
  - microscopic appearance 27
  - relationship to pernicious anaemia 65 66
  - physical examination in 36 37
  - psychic stress and 40
  - gastric secretion in 46 47 48
  - sex incidence 33 38
  - association with Sjogren's disease 39
  - structural changes in 24
  - symptomatology 33
  - tea drinking and 41
  - atrophy of tongue in 37
  - treatment with acid 33
  - treatment with alkali 51
  - treatment with iron 51
  - treatment with penicillin 51
  - relationship to peptic ulcer 45
  - Wits's anaemia and 48
  - causing vitamin B<sub>12</sub> deficiency 51
  - vitamin deficiency in 38
  - X-ray irradiation 37 40 44 45
- Gastritis diffuse giant hypertrophic 25 53
- — — barium meal in 54
  - — — biopsy in 54
  - — — differential diagnosis 54
- Gastritis diffuse giant hypertrophic relation to chronic gastritis 24 25 53
- — — gastroscopy in 54
  - — — macroscopic appearance 53
  - — — microscopic appearance 53
  - — — relation to neoplasm 53
  - — — treatment of 54
- Gastritis zonal 47
- Gastroscopy 5
- in acute gastritis 21 22
  - in alcoholism 39
  - complications 5
  - relation to gastric biopsy 5
  - findings in acute gastritis 5
  - findings in chronic gastritis 5
  - findings in diffuse giant hypertrophic gastritis 54
  - findings in pernicious anaemia 5
  - technique 5
  - instrument Benedict 7
  - instrument Cameron 5
  - instrument Hermon Taylor 5 7
  - instrument Schindler 5 7
- Giant hypertrophic gastritis *see* Gastritis diffuse giant hypertrophic
- Haematemesis
- after gastric biopsy 16 17
  - in acute gastritis 20
- Haemochromatosis 39
- Haemorrhage
- after gastric biopsy 16 17
  - in chronic gastritis 33 36 50
  - in diffuse giant hypertrophic gastritis 54
- Hepatomegaly and alcoholism 2
- Hypertrophic gastritis *see also* Gastritis diffuse giant hypertrophic
- Influenza acute gastritis in 23
- Intestinal metaplasia
- in gastric atrophy 57
  - in chronic gastritis 29
- Intrinsic factor of Castle 64 66 67
- in gastric mucoprotein 67
- Iron absorption 49 50

- Iron deficiency  
 relation to achlorhydria 48  
 in Wits anaemia 48 49 50
- Irradiation *see* \ ray irradiation
- Malnutrition *see* Diet
- Megaloblastic anaemia  
 caused by Epanutin 70  
 gastric biopsy in 15 63 70 71  
 gastric changes in 66 67 68
- Mental changes in vitamin B<sub>12</sub>  
 deficiency 6~
- Oesophagus perforation after  
 biopsy 17  
 bleeding from varices 36
- Pain in gastritis *see* Dyspepsia
- Pancreatitis -
- Pepsinogen  
 plasma 4  
 urinary 4
- Pepsin secretion *see also* Gastric  
 secretion  
 correlation with gastric biopsy  
 15  
 in peptic ulcer 45
- Peripheral neuritis 71  
 alcoholic simulating subacute  
 combined degeneration of the  
 cord 71
- Pernicious anaemia *see also* Vita-  
 min B<sub>12</sub>  
 aetiology of 64 65 66  
 antibiotics in 70  
 relation to cancer 66  
 in childhood 68 69  
 definition of 69  
 dyspepsia in 61 6~  
 familial incidence of 65 66 68  
 69  
 gastric cytology in 60  
 gastroscopy in 5  
 preceded by gastric atrophy 61  
 normal gastric biopsy in 68  
 and gastric changes 57 59 66  
 history 57  
 intrinsic factor in 67  
 macroscopic appearance of the  
 stomach in 57  
 microscopic appearance of the  
 stomach in 57 58  
 oral administration of vitamin  
 B<sub>12</sub> 70
- Pernicious anaemia—*contd*  
 subacute combined degeneration  
 of the cord in the absence of  
 6~ 63  
 symptomatology 61  
 tongue in 61  
 transfusion in 70  
 treatment of 69 70
- Physical examination 2 3  
 in chronic gastritis 36 37
- Pneumonia acute gastritis in 23
- Psychic stress cause of gastritis 42
- Radiation *see also* \ ray irradiation  
 and Gastritis chronic
- Radiology in diffuse giant hyper-  
 trophic gastritis 4 54  
 radiology in chronic gastritis 4
- Rugae  
 effect of distension 25  
 in chronic gastritis 25  
 in diffuse giant hypertrophic  
 gastritis 54  
 pronounced 4
- Salmonella infections 20
- Sjogren's disease 39
- Staining of gastric biopsy section  
 13
- Staphylococcal  
 food poisoning 21 23 39  
 gastroenteritis 21 39
- Subacute combined degeneration  
 of the cord 57 59  
 achlorhydria in 59  
 gastric biopsy indicated in 15  
 diagnosis with gastric biopsy  
 71  
 relation to gastric changes 66  
 macroscopic appearance of the  
 stomach in 57 58  
 microscopic appearance of the  
 stomach in 59  
 peripheral neuritis 3  
 development without pernicious  
 anaemia 6~ 63  
 symptomatology 61  
 treatment of 69 70  
 cobalt labelled vitamin B<sub>12</sub>  
 absorption test in 63  
 relation to vitamin B<sub>12</sub> deficiency  
 68
- Superficial gastritis *see* Gastritis  
 chronic

- Tea cause of gastritis 41
- Test meal  
correlation with gastric biopsy 3 4  
histamine 3 66  
insulin 4  
maximum secretory response 4  
overnight 4  
tubeless 4
- Tongue  
atrophy in chronic gastritis 37  
in pernicious anaemia 61
- Trichrome stain for gastric biopsy 11, 12
- Ulcer 2 3  
age of onset 36  
bleeding from 36  
duodenal 42 44 45 46  
gastric secretion in 45 46  
relationship of chronic gastritis to 45  
zonal gastritis in relation to 45 47
- Vitamin B<sub>12</sub> *see also* Pernicious anaemia  
mode of action 67 68  
<sup>58</sup>cobalt labelled in absorption test 15 63 71  
deficiency following total gastrectomy 62 68  
deficiency manifestations of 61 62 63  
in gastric atrophy 61
- Vitamin B<sub>12</sub>—*contd*  
non reversibility of gastric atrophy to 15 60 70  
deficiency due to chronic gastritis 51  
relation to intrinsic factor of Castle 67 68  
storage in liver 68  
mental changes in 62  
oral administration of 70  
in pernicious anaemia 15  
serum estimations 71  
relation to subacute combined degeneration of the cord 68  
transport in serum 67  
in treatment of pernicious anaemia 69 70  
in treatment of subacute combined degeneration of the cord 69 70  
absorption in Witts anaemia 46 49
- Vitamin deficiency  
in alcoholism 41  
causing cardiac enlargement 3  
in chronic gastritis 38  
causing peripheral neuritis 2  
causing tachycardia 3
- Witts anaemia *see* Anaemia
- X ray irradiation 42 44  
as a carcinogen 49  
effect on gastric secretion 42 44  
cause of gastritis 37 42 44 45

